



HoxD4 (P-18): sc-82928

BACKGROUND

The Hox proteins are a family of transcription factors that play a role in development and cellular differentiation by regulating downstream target genes. Specifically, the Hox proteins direct DNA-protein and protein-protein interactions that assist in determining the morphologic features associated with the anterior-posterior body axis. Hox proteins are involved in controlling axial patterning, leukemias and hereditary malformations. HoxD4 (homeobox protein D4), also known as HOX4B, is a 255 amino acid protein that localizes to the nucleus and contains one homeobox DNA-binding domain. One of several members of the homeobox superfamily, HoxD4 functions as a sequence-specific transcription factor that is important for the correct positioning of developing limb buds on the anterior-posterior axis. Mutations in the gene encoding HoxD4 are associated with the pathogenesis of childhood acute lymphoblastic leukemia.

REFERENCES

- Mavilio, F., Simeone, A., Giampaolo, A., Faiella, A., Zappavigna, V., Acampora, D., Poiana, G., Russo, G., Peschle, C. and Boncinelli, E. 1986. Differential and stage-related expression in embryonic tissues of a new human homeobox gene. *Nature* 324: 664-668.
- Oliver, G., Sidell, N., Fiske, W., Heinzmann, C., Mohandas, T., Sparkes, R.S. and De Robertis, E.M. 1989. Complementary homeo protein gradients in developing limb buds. *Genes Dev.* 3: 641-650.
- Kim, Y.H., Choi, C.Y., Lee, S.J., Conti, M.A. and Kim, Y. 1998. Homeodomain-interacting protein kinases, a novel family of corepressors for homeodomain transcription factors. *J. Biol. Chem.* 273: 25875-25879.
- Del Campo, M., Jones, M.C., Veraksa, A.N., Curry, C.J., Jones, K.L., Mascarello, J.T., Ali-Kahn-Catts, Z., Drumheller, T. and McGinnis, W. 1999. Monodactylous limbs and abnormal genitalia are associated with hemizygosity for the human 2q31 region that includes the HOXD cluster. *Am. J. Hum. Genet.* 65: 104-110.
- Zákány, J. and Duboule, D. 1999. Hox genes and the making of sphincters. *Nature* 401: 761-762.
- Shen, W.F., Krishnan, K., Lawrence, H.J. and Largman, C. 2001. The Hox homeodomain proteins block CBP histone acetyltransferase activity. *Mol. Cell. Biol.* 21: 7509-7522.
- Kosaki, K., Kosaki, R., Suzuki, T., Yoshihashi, H., Takahashi, T., Sasaki, K., Tomita, M., McGinnis, W. and Matsuo, N. 2002. Complete mutation analysis panel of the 39 human HOX genes. *Teratology* 65: 50-62.
- van Scherpenzeel Thim, V., Rémacle, S., Picard, J., Cornu, G., Gofflot, F., Rezsóhazy, R. and Verellen-Dumoulin, C. 2005. Mutation analysis of the HOX paralogous 4-13 genes in children with acute lymphoid malignancies: identification of a novel germline mutation of HOXD4 leading to a partial loss-of-function. *Hum. Mutat.* 25: 384-395.

CHROMOSOMAL LOCATION

Genetic locus: HOXD4 (human) mapping to 2q31.1; Hoxd4 (mouse) mapping to 2 C3.

SOURCE

HoxD4 (P-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HoxD4 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-82928 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-82928 X, 200 µg/0.1 ml.

APPLICATIONS

HoxD4 (P-18) is recommended for detection of HoxD4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Hox family members.

Suitable for use as control antibody for HoxD4 siRNA (h): sc-75295, HoxD4 siRNA (m): sc-75296, HoxD4 shRNA Plasmid (h): sc-75295-SH, HoxD4 shRNA Plasmid (m): sc-75296-SH, HoxD4 shRNA (h) Lentiviral Particles: sc-75295-V and HoxD4 shRNA (m) Lentiviral Particles: sc-75296-V.

HoxD4 (P-18) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of HoxD4: 28 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.