

PARP-11 (L-17): sc-82967

BACKGROUND

Poly(ADP-ribosylation) is a method of DNA damage-dependent posttranslational modification that helps to rescue injured proliferating cells from cell death. The PARP (poly(ADP-ribose) polymerase) proteins comprise a superfamily of enzymes that functionally modify histones and other nuclear proteins, thereby preventing cell death. PARPs use NAD⁺ as a substrate to catalytically transfer ADP-ribose residues onto protein acceptors; a process that, when repeated multiple times, leads to the formation of poly(ADP-ribose) chains on the protein. The presence of these chains alters the function of the target protein and promotes cell survival. PARP proteins are implicated in a variety of diseases, including cancer, neurodegenerative and inflammatory disorders.

REFERENCES

- Hans, M.A., et al. 1999. Overexpression of dominant negative PARP interferes with tumor formation of HeLa cells in nude mice: evidence for increased tumor cell apoptosis *in vivo*. *Oncogene* 18: 7010-7015.
- Aguiar, R.C., et al. 2005. B-aggressive lymphoma family proteins have unique domains that modulate transcription and exhibit poly(ADP-ribose) polymerase activity. *J. Biol. Chem.* 280: 33756-33765.
- Chou, H.Y., et al. 2006. CDK-dependent activation of poly(ADP-ribose) polymerase member 10 (PARP-10). *J. Biol. Chem.* 281: 15201-15207.
- Goenka, S., et al. 2007. Collaborator of Stat6 (CoaSt6)-associated poly(ADP-ribose) polymerase activity modulates Stat6-dependent gene transcription. *J. Biol. Chem.* 282: 18732-18739.
- Liu, X., et al. 2008. Poly (ADP-ribose) polymerase activity regulates apoptosis in HeLa cells after alkylating DNA damage. *Cancer Biol. Ther.* 7: 934-941.
- Elser, M., et al. 2008. Poly(ADP-ribose) polymerase 1 promotes tumor cell survival by coactivating hypoxia-inducible factor-1-dependent gene expression. *Mol. Cancer Res.* 6: 282-290.

CHROMOSOMAL LOCATION

Genetic locus: PARP11 (human) mapping to 12p13.32; Parp11 (mouse) mapping to 6 F3.

SOURCE

PARP-11 (L-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PARP-11 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-82967 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-82967 X, 200 µg/0.1 ml.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

PARP-11 (L-17) is recommended for detection of PARP-11 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other PARP family members.

PARP-11 (L-17) is also recommended for detection of PARP-11 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for PARP-11 siRNA (h): sc-76052, PARP-11 siRNA (m): sc-76053, PARP-11 shRNA Plasmid (h): sc-76052-SH, PARP-11 shRNA Plasmid (m): sc-76053-SH, PARP-11 shRNA (h) Lentiviral Particles: sc-76052-V and PARP-11 shRNA (m) Lentiviral Particles: sc-76053-V.

PARP-11 (L-17) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of PARP-11 isoforms: 39/32/29 kDa.

Positive Controls: PARP-11 (m): 293T Lysate: sc-122383 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



PARP-11 (L-17): sc-82967. Western blot analysis of PARP-11 expression in non-transfected 293T: sc-117752 (A), mouse PARP-11 transfected 293T: sc-122383 (B) and HeLa (C) whole cell lysates.



PARP-11 (L-17): sc-82967. Immunofluorescence staining of methanol-fixed HeLa cells showing membrane localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.