

RYK (H-190): sc-83081

BACKGROUND

RYK (related to receptor tyrosine kinase) is a chemorepulsive axon guidance receptor required for establishing major axon tracts, such as the corticospinal tract and the corpus callosum, in the developing nervous system. RYK belongs to the RTK family and is ubiquitously expressed in both developing and adult tissues. Compared to other RTK family members, RYK has a relatively short extracellular domain. In addition, RYK contains two N-terminal leucine-rich motifs that resemble the N-terminal domain of WIF-1. RYK functions as a Wnt receptor and produces a repulsive guidance signal upon binding of Wnt-1, 3, 3a or 5a to its WIF-like domain. The loss of functional RYK results in the formation of contralateral axon bundles and the disruption of cardiac, skeletal and craniofacial development, ultimately resulting in perinatal death. RYK is also strongly expressed in malignant ovarian tumors and may play a role in tumorigenesis.

REFERENCES

- Masiakowski, P. and Carroll, R.D. 1992. A novel family of cell surface receptors with tyrosine kinase-like domain. *J. Biol. Chem.* 267: 26181-26190.
- Paganoni, S. and Ferreira, A. 2003. Expression and subcellular localization of Ror tyrosine kinase receptors are developmentally regulated in cultured hippocampal neurons. *J. Neurosci. Res.* 73: 429-440.
- Yoda, A., et al. 2003. Expression and function of the Ror-family receptor tyrosine kinases during development: lessons from genetic analyses of nematodes, mice, and humans. *J. Recept. Signal Transduct. Res.* 23: 1-15.
- Shabani, M., et al. 2007. Overexpression of orphan receptor tyrosine kinase ROR1 as a putative tumor-associated antigen in Iranian patients with acute lymphoblastic leukemia. *Tumour Biol.* 28: 318-326.
- Winkel, A., et al. 2008. Wnt-ligand-dependent interaction of TAK1 (TGF β -activated kinase-1) with the receptor tyrosine kinase ROR2 modulates canonical Wnt-signalling. *Cell. Signal.* 20: 2134-2144.
- Baskar, S., et al. 2008. Unique cell surface expression of receptor tyrosine kinase ROR1 in human B cell chronic lymphocytic leukemia. *Clin. Cancer Res.* 14: 396-404.
- Daneshmanesh, A.H., et al. 2008. ROR1, a cell surface receptor tyrosine kinase is expressed in chronic lymphocytic leukemia and may serve as a putative target for therapy. *Int. J. Cancer* 123: 1190-1195.
- Shabani, M., et al. 2008. Expression profile of orphan receptor tyrosine kinase (ROR1) and Wilms' tumor gene 1 (WT1) in different subsets of B cell acute lymphoblastic leukemia. *Leuk. Lymphoma* 49: 1360-1367.

CHROMOSOMAL LOCATION

Genetic locus: RYK (human) mapping to 3q22.2; RYK (mouse) mapping to 9 F1.

SOURCE

RYK (H-190) is a rabbit polyclonal antibody raised against amino acids 48-220 mapping within an extracellular domain of RYK of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

RYK (H-190) is recommended for detection of RYK of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RYK siRNA (h): sc-72221, RYK siRNA (m): sc-72222, RYK shRNA Plasmid (h): sc-72221-SH, RYK shRNA Plasmid (m): sc-72222-SH, RYK shRNA (h) Lentiviral Particles: sc-72221-V and RYK shRNA (m) Lentiviral Particles: sc-72222-V.

Molecular Weight of RYK: 90 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfaction
 Guaranteed

Try **RYK (F35 P7 D7 F5): sc-83082**, our highly recommended monoclonal alternative to RYK (H-190).