

SETD4 (P-15): sc-83750

BACKGROUND

SETD4 (SET domain-containing protein 4) is a 440 amino acid protein that contains one SET domain and is expressed as three isoforms produced by alternative splicing. The gene that encodes SETD4 maps to human chromosome 21. The smallest of the human chromosomes, 21 makes up about 1.5% of the human genome. Chromosome 21 contains nearly 300 genes and 47 million base pairs. Down syndrome, also known as trisomy 21, is the disease most commonly associated with chromosome 21. Alzheimer's disease, Jervell and Lange-Nielsen syndrome and amyotrophic lateral sclerosis are also associated with chromosome 21. Translocations are found to occur between chromosome 21 and 8, and chromosome 21 and 12, in certain leukemias.

REFERENCES

1. Tesson, F., et al. 1996. Exclusion of KCNE1 (IsK) as a candidate gene for Jervell and Lange-Nielsen syndrome. *J. Mol. Cell. Cardiol.* 28: 2051-2055.
2. Tyson, J., et al. 1997. IsK and KvLQT1: mutation in either of the two subunits of the slow component of the delayed rectifier potassium channel can cause Jervell and Lange-Nielsen syndrome. *Hum. Mol. Genet.* 6: 2179-2185.

CHROMOSOMAL LOCATION

Genetic locus: SETD4 (human) mapping to 21q22.12; Setd4 (mouse) mapping to 16 C4.

SOURCE

SETD4 (P-15) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of SETD4 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-83750 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SETD4 (P-15) is recommended for detection of SETD4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with isoforms A, B or 3.

SETD4 (P-15) is also recommended for detection of SETD4 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for SETD4 siRNA (h): sc-91446, SETD4 siRNA (m): sc-153385, SETD4 shRNA Plasmid (h): sc-91446-SH, SETD4 shRNA Plasmid (m): sc-153385-SH, SETD4 shRNA (h) Lentiviral Particles: sc-91446-V and SETD4 shRNA (m) Lentiviral Particles: sc-153385-V.


Molecular Weight of SETD4: 44 kDa.

Positive Controls: HL-60 whole cell lysate: sc-2209, K-562 whole cell lysate: sc-2203 or mouse testis extract: sc-2405.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



SETD4 (P-15): sc-83750. Western blot analysis of SETD4 expression in HEK293 (A), K-562 (B) and HL-60 (C) whole cell lysates and rat cerebellum (D) and mouse testis (E) tissue extracts.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **SETD4 (F-3): sc-514060**, our highly recommended monoclonal alternative to SETD4 (P-15).