

Cdc42 (B-8): sc-8401

BACKGROUND

The superfamily of GTP-binding proteins, for which the Ras proteins are prototypes, has been implicated in regulation of diverse biological activities involving various aspects of cell growth and division. One mammalian member of the family, Cdc42, has an amino acid sequence that is similar to those of various members of the Ras superfamily proteins, including N-, K- and H-Ras, Rho proteins and the Rac proteins. On the basis of *in vitro* phosphorylation studies, it has been suggested that human Cdc42 may function in the signaling pathway of the EGF receptor or related growth factor receptor protein kinases. The Dbl oncogene has been shown to specifically catalyze dissociation of GDP from human Cdc42.

CHROMOSOMAL LOCATION

Genetic locus: CDC42 (human) mapping to 1p36.12; Cdc42 (mouse) mapping to 4 D3.

SOURCE

Cdc42 (B-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 150-182 at the C-terminus of Cdc42 of human origin.

PRODUCT

Each vial contains 200 µg IgG₃ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Cdc42 (B-8) is available conjugated to agarose (sc-8401 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-8401 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; and to either phycoerythrin (sc-8401 PE), fluorescein (sc-8401 FITC), Alexa Fluor® 488 (sc-8401 AF488) or Alexa Fluor® 647 (sc-8401 AF647), 200 µg/ml, for IF, IHC(P) and FCM.

Blocking peptide available for competition studies, sc-8401 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

Cdc42 (B-8) is recommended for detection of Cdc42 of mouse, rat, human, *Drosophila melanogaster* and *Xenopus laevis* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Cdc42 (B-8) is also recommended for detection of Cdc42 in additional species, including equine, canine, bovine, porcine and avian.


Suitable for use as control antibody for Cdc42 siRNA (h): sc-29256, Cdc42 siRNA (m): sc-29257, Cdc42 shRNA Plasmid (h): sc-29256-SH, Cdc42 shRNA Plasmid (m): sc-29257-SH, Cdc42 shRNA (h) Lentiviral Particles: sc-29256-V and Cdc42 shRNA (m) Lentiviral Particles: sc-29257-V.

Molecular Weight of Cdc42: 25 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



Cdc42 (B-8): sc-8401. Near-infrared western blot analysis of Cdc42 expression in Jurkat (A), 3611-RF (B), HeLa (C), THP-1 (D), A-431 (E) and Hep G2 (F) whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgGκ BP-CFL 680: sc-516180.

Cdc42 (B-8): sc-8401. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic staining of myocytes (B).

SELECT PRODUCT CITATIONS

- Sahai, E., et al. 2001. Cross-talk between Ras and Rho signalling pathways in transformation favours proliferation and increased motility. *EMBO J.* 20: 755-766.
- Foger, N., et al. 2001. Involvement of CD44 in cytoskeleton rearrangement and raft reorganization in T cells. *J. Cell Sci.* 114: 1169-1178.
- Ko, H.K., et al. 2014. Suppression of chemotaxis by SSeCKS via scaffolding of phosphoinositol phosphates and the recruitment of the Cdc42 GEF, Frabin, to the leading edge. *PLoS ONE* 9: e111534.
- Ruppersburg, C.C., et al. 2014. The Ca²⁺-activated Cl⁻ channel ANO1/TMEM16A regulates primary ciliogenesis. *Mol. Biol. Cell* 25: 1793-1807.
- Seo, H.S., et al. 2015. Identification of biomarkers regulated by rexinoids (LGD1069, LG100268 and Ro25-7386) in human breast cells using Affymetrix microarray. *Mol. Med. Rep.* 12: 800-818.
- Oprea, T.I., et al. 2015. Novel activities of select NSAID R-Enantiomers against Rac1 and Cdc42 GTPases. *PLoS ONE* 10: e0142182.
- Abraham, S., et al. 2015. A Rac/Cdc42 exchange factor complex promotes formation of lateral filopodia and blood vessel lumen morphogenesis. *Nat. Commun.* 6: 7286.
- Miller, G.J., et al. 2015. Mechanical microenvironments and protein expression associated with formation of different skeletal tissues during bone healing. *Biomech. Model. Mechanobiol.* 14: 1239-1253.
- Fan, S.H., et al. 2016. Endosomal Na⁺/H⁺ exchanger NHE5 influences MET recycling and cell migration. *Mol. Biol. Cell* 27: 702-715.

RESEARCH USE

For research use only, not for use in diagnostic procedures.