

## ESD (T-20): sc-84107

### BACKGROUND

ESD (esterase D) is also known as S-formylglutathione hydrolase and is a 282 amino acid protein that is a member of the esterase D family. ESD is highly expressed in placenta, kidney, liver and erythrocytes, and is localized to the cytoplasm, as well as to cytoplasmic vesicles. The main function of ESD is to detoxify formaldehyde while providing energy. Formaldehyde is oxidized by ADH5 which yields S-formylglutathione. ESD then catalyzes the hydrolysis of S-formylglutathione to the reduced forms of formic acid and glutathione. In addition, ESD hydrolyzes a variety of different neutral ester substrates and can act as a carboxylesterase. ESD may also act as a cysteine hydrolase which is inactivated by thiol alkylating agents. ESD gene polymorphism can lead to reduced enzymatic activity which may cause susceptibility to many conditions, including toxic liver cirrhosis, retinoblastoma, obesity and autism.

### REFERENCES

- Harms, N., et al. 1996. S-formylglutathione hydrolase of *Paracoccus denitrificans* is homologous to human esterase D: a universal pathway for formaldehyde detoxification? *J. Bacteriol.* 178: 6296-6299.
- McAuley, K.E., et al. 2003. Purification, crystallization and preliminary X-ray diffraction analysis of S-formylglutathione hydrolase from *Arabidopsis thaliana*: effects of pressure and selenomethionine substitution on space-group changes. *Acta Crystallogr. D Biol. Crystallogr.* 59: 2272-2274.
- Yurimoto, H., et al. 2003. Physiological role of S-formylglutathione hydrolase in C<sub>1</sub> metabolism of the methylotrophic yeast *Candida boidinii*. *Microbiology* 149: 1971-1979.
- Yuasa, I., et al. 2004. Molecular basis of ESD\*5 and ESD\*7 and haplotype analysis with new polymorphisms in introns. *Hum. Biol.* 76: 479-488.
- Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 133280. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Gonzalez, C.F., et al. 2006. Molecular basis of formaldehyde detoxification. Characterization of two S-formylglutathione hydrolases from *Escherichia coli*, FrmB and YeiG. *J. Biol. Chem.* 281: 14514-14522.
- Cummins, I., et al. 2006. Unique regulation of the active site of the serine esterase S-formylglutathione hydrolase. *J. Mol. Biol.* 359: 422-432.
- Legler, P.M., et al. 2008. Structural characterization and reversal of the natural organophosphate resistance of a D-type esterase, *Saccharomyces cerevisiae* S-formylglutathione hydrolase. *Biochemistry* 47: 9592-9601.
- Okunuki, Y., et al. 2008. Proteomic surveillance of retinal autoantigens in endogenous uveitis: implication of esterase D and brain-type creatine kinase as novel autoantigens. *Mol. Vis.* 14: 1094-1104.

### CHROMOSOMAL LOCATION

Genetic locus: ESD (human) mapping to 13q14.2; ESD (mouse) mapping to 14 D3.

### SOURCE

ESD (T-20) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of ESD of human origin.

### PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-84107 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

ESD (T-20) is recommended for detection of ESD of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ESD (T-20) is also recommended for detection of ESD in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ESD siRNA (h): sc-105338, ESD siRNA (m): sc-144943, ESD shRNA Plasmid (h): sc-105338-SH, ESD shRNA Plasmid (m): sc-144943-SH, ESD shRNA (h) Lentiviral Particles: sc-105338-V and ESD shRNA (m) Lentiviral Particles: sc-144943-V.

Molecular Weight of ESD: 31 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.