

MTRF1 (C-15): sc-84213



The Power to Question

BACKGROUND

MTRF1 (mitochondrial translational release factor 1), also known as RF1 or MRF1, is a 445 amino acid protein that localizes to the mitochondrion and belongs to the prokaryotic/mitochondrial release factor family of chain-terminating proteins. Functioning in a similar manner to related yeast and bacterial proteins, MTRF1 acts as a peptide chain release factor that directs translational termination, specifically in response to the stop codons AGG and AGA. The gene encoding MTRF1 maps to human chromosome 13, which houses over 400 genes, such as BRCA2 and RB1, and comprises nearly 4% of the human genome. Trisomy 13, also known as Patau syndrome, is deadly and the few who survive past one year suffer from permanent neurologic defects, difficulty eating and vulnerability to serious respiratory infections.

REFERENCES

- Zhang, Y. and Spremulli, L.L. 1998. Identification and cloning of human mitochondrial translational release factor 1 and the ribosome recycling factor. *Biochim. Biophys. Acta* 1443: 245-250.
- Hansen, L.L., Jorgensen, R. and Justesen, J. 2000. Assignment of the human mitochondrial translational release factor 1 (MTRF1) to chromosome 13q14.1→q14.3 and of the human mitochondrial ribosome recycling factor (MRRF) to chromosome 9q32→q34.1 with radiation hybrid mapping. *Cytogenet. Cell Genet.* 88: 91-92.
- Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 604601. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Lekomtsev, S.A. 2007. Non-standard genetic codes and translation termination. *Mol. Biol.* 41: 964-972.
- Soleimanpour-Lichaei, H.R., Kühl, I., Gaisne, M., Passos, J.F., Wydro, M., Rorbach, J., Temperley, R., Bonnefoy, N., Tate, W., Lightowlers, R. and Chrzanowska-Lightowlers, Z. 2007. mtRF1a is a human mitochondrial translation release factor decoding the major termination codons UAA and UAG. *Mol. Cell.* 27: 745-757.
- Kutner, J. 2007. Termination of prokaryotic and eukaryotic translation. *Postepy Biochem.* 53: 420-430.
- Ishizawa, T., Nozaki, Y., Ueda, T. and Takeuchi, N. 2008. The human mitochondrial translation release factor HMRF1L is methylated in the GGQ motif by the methyltransferase HMPmC. *Biochem. Biophys. Res. Commun.* 373: 99-103.
- Nozaki, Y., Matsunaga, N., Ishizawa, T., Ueda, T. and Takeuchi, N. 2008. HMRF1L is a human mitochondrial translation release factor involved in the decoding of the termination codons UAA and UAG. *Genes Cells.* 13: 429-438.

CHROMOSOMAL LOCATION

Genetic locus: MTRF1 (human) mapping to 13q14.11; Mtrf1 (mouse) mapping to 14 D3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

MTRF1 (C-15) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the C-terminus of MTRF1 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-84213 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MTRF1 (C-15) is recommended for detection of MTRF1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MTRF1 siRNA (h): sc-106261, MTRF1 siRNA (m): sc-149692, MTRF1 shRNA Plasmid (h): sc-106261-SH, MTRF1 shRNA Plasmid (m): sc-149692-SH, MTRF1 shRNA (h) Lentiviral Particles: sc-106261-V and MTRF1 shRNA (m) Lentiviral Particles: sc-149692-V.

Molecular Weight of MTRF1: 52 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.