

E-cadherin (G-10): sc-8426

BACKGROUND

E-cadherin Antibody (G-10) is a high quality monoclonal E-cadherin antibody (also designated CD324 antibody, ECAD antibody or CDH1 antibody) suitable for the detection of the E-cadherin protein of mouse, rat and human origin. E-cadherin Antibody (G-10) is available as both the non-conjugated anti-E-cadherin antibody form, as well as multiple conjugated forms of anti-E-cadherin antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor[®] conjugates. Cadherins comprise a family of Ca²⁺-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. Members of this family of adhesion proteins include rat cadherin K (and its human homolog, cadherin-6), R-cadherin, B-cadherin, E/P cadherin and cadherin-5. The classical cadherins, E-, N- and P-cadherin, consist of large extracellular domains characterized by a series of five homologous NH2 terminal repeats. The most distal of these cadherins is thought to be responsible for binding specificity, transmembrane domains and carboxy terminal intracellular domains. The relatively short intracellular domains interact with a variety of cytoplasmic proteins, such as β -catenin, to regulate cadherin function.

CHROMOSOMAL LOCATION

Genetic locus: CDH1 (human) mapping to 16q22.1; Cdh1 (mouse) mapping to 8 D3.

SOURCE

E-cadherin (G-10) is a mouse monoclonal antibody raised against amino acids 600-707 mapping within an extracellular domain of E-cadherin of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

E-cadherin (G-10) is available conjugated to agarose (sc-8426 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-8426 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-8426 PE), fluorescein (sc-8426 FITC), Alexa Fluor[®] 488 (sc-8426 AF488), Alexa Fluor[®] 546 (sc-8426 AF546), Alexa Fluor[®] 594 (sc-8426 AF594) or Alexa Fluor[®] 647 (sc-8426 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-8426 AF680) or Alexa Fluor[®] 790 (sc-8426 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS


E-cadherin (G-10) is recommended for detection of E-cadherin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for E-cadherin siRNA (h): sc-35242, E-cadherin siRNA (m): sc-35243, E-cadherin shRNA Plasmid (h): sc-35242-SH, E-cadherin shRNA Plasmid (m): sc-35243-SH, E-cadherin shRNA (h) Lentiviral Particles: sc-35242-V and E-cadherin shRNA (m) Lentiviral Particles: sc-35243-V.


Molecular Weight of mature E-cadherin: 120/80 kDa.

Molecular Weight of E-cadherin precursor: 135 kDa.

DATA



E-cadherin (G-10) Alexa Fluor[®] 488: sc-8426 AF488. Direct fluorescent western blot analysis of E-cadherin expression in LNCaP (A), MCF7 (B) and Caco-2 (C) whole cell lysates. Blocked with UltraCruz[®] Blocking Reagent: sc-516214. Cruz Marker[™] Molecular Weight Standards detected with Cruz Marker[™] MW Tag-Alexa Fluor[®] 647: sc-516791.



E-cadherin (G-10): sc-8426. Immunofluorescence staining of methanol-fixed LNCaP cells showing membrane staining (A). Immunoperoxidase staining of formalin-fixed, paraffin-embedded normal human tonsil showing membrane localization (B).

SELECT PRODUCT CITATIONS

1. Lipecka, J., et al. 2002. Distribution of ClC-2 chloride channel in rat and human epithelial tissues. *Am. J. Physiol., Cell Physiol.* 282: C805-C816.
2. Tsogbadrakh, B., et al. 2018. HL156A, a novel pharmacological agent with potent adenosine-monophosphate-activated protein kinase (AMPK) activator activity ameliorates renal fibrosis in a rat unilateral ureteral obstruction model. *PLoS ONE* 13: e0201692.
3. Sengez, B., et al. 2019. The transcription factor Elf3 is essential for a successful mesenchymal to epithelial transition. *Cells* 8: 858.
4. Karagonlar, Z.F., et al. 2020. A novel function for KLF4 in modulating the de-differentiation of EpCAM⁺/CD133⁺ nonStem cells into EpCAM⁺/CD133⁺ liver cancer stem cells in HCC cell line HuH7. *Cells* 9: 1198.
5. Wijshake, T., et al. 2021. Tumor-suppressor function of Beclin 1 in breast cancer cells requires E-cadherin. *Proc. Natl. Acad. Sci. USA* 118: e2020478118.

RESEARCH USE

For research use only, not for use in diagnostic procedures.