


Actin (C-2): sc-8432



The Power to Question

BACKGROUND

All eukaryotic cells express Actin, which often constitutes as much as 50% of total cellular protein. Actin filaments can form both stable and labile structures and are crucial components of microvilli and the contractile apparatus of muscle cells. While lower eukaryotes, such as yeast, have only one Actin gene, higher eukaryotes have several isoforms encoded by a family of genes. At least six types of Actin are present in mammalian tissues and fall into three classes. α -Actin expression is limited to various types of muscle, whereas β -Actin and γ -Actin are the principle constituents of filaments in other tissues. Members of the small GTPase family regulate the organization of the Actin cytoskeleton. Rho controls the assembly of Actin stress fibers and focal adhesion. Rac regulates Actin filament accumulation at the plasma membrane. Cdc42 stimulates formation of filopodia.

REFERENCES

- Doolittle, R.F. 1995. The origins and evolution of eukaryotic proteins. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 349: 235-240.
- Maccioni, R.B. and Cambiazo, V. 1995. Role of microtubule-associated proteins in the control of microtubule assembly. *Physiol. Rev.* 75: 835-864.
- Schutt, C.E., et al. 1995. A discourse on modeling F-Actin. *J. Struct. Biol.* 115: 186-198.
- Barkalow, K. and Hartwig, J.H. 1995. Actin cytoskeleton. Setting the pace of cell movement. *Curr. Biol.* 5: 1000-1002.

SOURCE

Actin (C-2) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 350-375 at the C-terminus of Actin of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Actin (C-2) is available conjugated to agarose (sc-8432 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-8432 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-8432 PE), fluorescein (sc-8432 FITC), Alexa Fluor[®] 488 (sc-8432 AF488), Alexa Fluor[®] 546 (sc-8432 AF546), Alexa Fluor[®] 594 (sc-8432 AF594) or Alexa Fluor[®] 647 (sc-8432 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-8432 AF680) or Alexa Fluor[®] 790 (sc-8432 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

In addition, Actin (C-2) is available conjugated to either TRITC (sc-8432 TRITC, 200 μ g/ml), Alexa Fluor[®] 405 (sc-8432 AF405, 200 μ g/ml) or PerCP (sc-8432 PerCP), 100 tests in 2 ml, for IF, IHC(P) and FCM.

Blocking peptide available for competition studies, sc-8432 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Actin (C-2) is recommended for detection of a broad range of Actin isoforms of mouse, rat, human and *Xenopus laevis* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500), flow cytometry (1 μ g per 1 x 10⁶ cells) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Actin (C-2) is also recommended for detection of a broad range of Actin isoforms in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Actin siRNA (h): sc-29191, Actin siRNA (m): sc-29192, Actin shRNA Plasmid (h): sc-29191-SH, Actin shRNA Plasmid (m): sc-29192-SH, Actin shRNA (h) Lentiviral Particles: sc-29191-V and Actin shRNA (m) Lentiviral Particles: sc-29192-V.


Molecular Weight of Actin: 43 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, MCF7 whole cell lysate: sc-2206 or CCRF-CEM cell lysate: sc-2225.

DATA



Actin (C-2): sc-8432. Western blot analysis of Actin expression in Jurkat (A), MCF7 (B), ZR-75-1 (C), CCRF-CEM (D) and HeLa (E) whole cell lysates and human colon tissue extract (F). Detection reagent used: m-IgG₁ BP-HRP: sc-525408.



Actin (C-2): sc-8432. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic staining of myocytes (A). Immunofluorescence staining of methanol-fixed HeLa cells showing cytoskeletal localization (B).

SELECT PRODUCT CITATIONS

- Matsushima, H. and Bogenmann, E. 1993. Stromal platelet-derived growth factor receptor α (PDGFR α) provides a therapeutic target independent of tumor cell PDGFR α expression in lung cancer xenografts. *Mol. Cell. Biol.* 13: 7447-7456.
- Zeni, C., et al. 2024. Formin binding protein 1 (FBNP1) regulates non-canonical Wnt signaling and vertebrate gastrulation. *Dev. Biol.* 515: 18-29.
- Seo, J., et al. 2025. Giant *Centella asiatica*, a novel cultivar rich in madecassoside and asiaticoside, suppresses α -melanocyte-stimulating hormone-induced melanogenesis through MC1R binding. *Int. J. Mol. Med.* 55: 13.

RESEARCH USE

For research use only, not for use in diagnostic procedures.