

AFG3L2 (L-16): sc-84687

BACKGROUND

AFG3L2 (ATPase family gene 3-like 2), also known as Paraplegin-like protein, is a multi-pass membrane metalloprotease that contains one AAA (ATPase associated with diverse cellular activities) domain, a zinc-dependent binding motif, an RNA-binding region and an ATP/GTP binding site. Localizing to the mitochondrial membrane, AFG3L2 is ubiquitously expressed with highest expression levels in skeletal muscle and heart. AFG3L2 shares 69% similarity with the yeast AFG3 protein and 49% similarity with Paraplegin, a protein of mitochondria that is thought to be involved in signal transduction and chaperone-like activities. In mitochondria, AFG3L2 forms a complex with Paraplegin that is believed to regulate essential protein quality control. Mutations in the gene encoding either one of these proteins can result in hereditary spastic paraplegia, a degenerative spinal cord disorder that is characterized by muscle spasms, stiffness in the legs and, in some cases, incontinence.

REFERENCES

- Banfi, S., Bassi, M.T., Andolfi, G., Marchitello, A., Zanotta, S., Ballabio, A., Casari, G. and Franco, B. 1999. Identification and characterization of AFG3L2, a novel paraplegin-related gene. *Genomics* 59: 51-58.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604581. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Olstad, O.K., Gautvik, V.T., Reppe, S., Rian, E., Jemtland, R., Ohlsson, C., Bruland, O.S. and Gautvik, K.M. 2003. Molecular heterogeneity in human osteosarcoma demonstrated by enriched mRNAs isolated by directional tag PCR subtraction cloning. *Anticancer Res.* 23: 2201-2216.
- Atorino, L., Silvestri, L., Koppen, M., Cassina, L., Ballabio, A., Marconi, R., Langer, T. and Casari, G. 2003. Loss of m-AAA protease in mitochondria causes complex I deficiency and increased sensitivity to oxidative stress in hereditary spastic paraplegia. *J. Cell Biol.* 163: 777-787.
- Nasir, J., Frima, N., Pickard, B., Malloy, M.P., Zhan, L. and Grünewald, R. 2006. Unbalanced whole arm translocation resulting in loss of 18p in dystonia. *Mov. Disord.* 21: 859-863.
- Duvezin-Caubet, S., Koppen, M., Wagener, J., Zick, M., Israel, L., Bernacchia, A., Jagasia, R., Rugarli, E.I., Imhof, A., Neupert, W., Langer, T. and Reichert, A.S. 2007. OPA1 processing reconstituted in yeast depends on the subunit composition of the m-AAA protease in mitochondria. *Mol. Biol. Cell* 18: 3582-3590.

CHROMOSOMAL LOCATION

Genetic locus: AFG3L2 (human) mapping to 18p11.21; Afg3l2 (mouse) mapping to 18 E1.

SOURCE

AFG3L2 (L-16) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an extracellular domain of AFG3L2 of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-84687 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

AFG3L2 (L-16) is recommended for detection of AFG3L2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

AFG3L2 (L-16) is also recommended for detection of AFG3L2 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for AFG3L2 siRNA (h): sc-72464, AFG3L2 siRNA (m): sc-140899, AFG3L2 shRNA Plasmid (h): sc-72464-SH, AFG3L2 shRNA Plasmid (m): sc-140899-SH, AFG3L2 shRNA (h) Lentiviral Particles: sc-72464-V and AFG3L2 shRNA (m) Lentiviral Particles: sc-140899-V.

Molecular Weight of full length AFG3L2: 80 kDa.


Molecular Weight of truncated AFG3L2: 65 kDa.

Positive Controls: rat skeletal muscle extract: sc-364810.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



AFG3L2 (L-16): sc-84687. Western blot analysis of AFG3L2 expression in rat skeletal muscle tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.