INO80C (E-20): sc-84749



The Power to Question

BACKGROUND

INO80C (INO80 complex subunit C), also known as C18orf37 or IES6, is a 192 amino acid protein that exists as multiple alternatively spliced isoforms and functions as a component of the multi-subunit chromatin-remodeling INO80 complex. The gene encoding INO80C maps to human chromosome 18q12.2, which houses over 300 protein-coding genes and contains nearly 76 million bases. There are a variety of diseases associated with defects in chromosome 18-localized genes, some of which include Trisomy 18 (also known as Edwards syndrome), Niemann-Pick disease, hereditary hemorrhagic telangiectasia, erythropoietic protoporphyria and follicular lymphomas.

REFERENCES

- Yoshikawa, T., et al. 1997. Isolation of chromosome 18-specific brain transcripts as positional candidates for bipolar disorder. Am. J. Med. Genet. 74: 140-149.
- Esterling, L.E., et al. 1997. An integrated physical map of 18p11.2: a susceptibility region for bipolar disorder. Mol. Psychiatry 2: 501-504.
- 3. Petek, E., et al. 2003. Characterisation of a 19-year-old "long-term survivor" with Edwards syndrome. Genet. Couns. 14: 239-244.
- 4. Jin, J., et al. 2005. A mammalian chromatin remodeling complex with similarities to the yeast INO80 complex. J. Biol. Chem. 280: 41207-41212.
- 5. Grosso, S., et al. 2005. Chromosome 18 aberrations and epilepsy: a review. Am. J. Med. Genet. A 134: 88-94.
- Nusbaum, C., et al. 2005. DNA sequence and analysis of human chromosome 18. Nature 437: 551-555.
- Pickard, B.S., et al. 2005. Candidate psychiatric illness genes identified in patients with pericentric inversions of chromosome 18. Psychiatr. Genet. 15: 37-44.
- 8. Farrants, A.K. 2008. Chromatin remodelling and actin organisation. FEBS Lett. 582: 2041-2050.
- 9. Conaway, R.C., et al. 2009. The INO80 chromatin remodeling complex in transcription, replication and repair. Trends Biochem. Sci. 34: 71-77.

CHROMOSOMAL LOCATION

Genetic locus: INO80C (human) mapping to 18q12.2; Ino80c (mouse) mapping to 18 A2.

SOURCE

INO80C (E-20) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of INO80C of human origin.

PRODUCT

Each vial contains 100 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-84749 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

INO80C (E-20) is recommended for detection of INO80C of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

INO80C (E-20) is also recommended for detection of INO80C in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for INO80C siRNA (h): sc-72684, INO80C siRNA (m): sc-142770, INO80C shRNA Plasmid (h): sc-72684-SH, INO80C shRNA Plasmid (m): sc-142770-SH, INO80C shRNA (h) Lentiviral Particles: sc-72684-V and INO80C shRNA (m) Lentiviral Particles: sc-142770-V.

Molecular Weight of INO80C: 21 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com