LOC284230 (Q-10): sc-84881



The Power to Question

BACKGROUND

The gene encoding human LOC284230 maps to chromosome 18, which houses over 300 protein-coding genes and contains nearly 76 million bases, representing about 2.5% of total DNA in cells. There are a variety of diseases associated with defects in chromosome 18-localized genes, some of which include Niemann-Pick disease, hereditary hemorrhagic telangiectasia, erythropoietic protoporphyria and follicular lymphomas. Also, three chromosomal abnormalities result from meiotic nondisjunction events of chromosome 18: monosomy 18p, trisomy 18 (also known as Edwards syndrome) and tetrasomy 18p. The LOC284230 gene product has been provisionally designated LOC284230 pending further characterization.

REFERENCES

- Grosso, S., Pucci, L., Di Bartolo, R.M., Gobbi, G., Bartalini, G., Anichini, C., Scarinci, R., Balestri, M., Farnetani, M.A., Cioni, M., Morgese, G. and Balestri, P. 2005. Chromosome 18 aberrations and epilepsy: a review. Am. J. Med. Genet. A 134A: 88-94.
- Stanyon, R., Rocchi, M., Capozzi, O., Roberto, R., Misceo, D., Ventura, M., Cardone, M.F., Bigoni, F. and Archidiacono, N. 2008. Primate chromosome evolution: ancestral karyotypes, marker order and neocentromeres. Chromosome Res. 16: 17-39.
- 3. Cerminara, C., Lo Castro, A., D'Argenzio, L., Galasso, C. and Curatolo, P. 2008. Epilepsy and deletion syndromes of chromosome 18: do not forget the short arm! Epilepsia 49: 1813-1814.
- Hollox, E.J., Barber, J.C., Brookes, A.J. and Armour, J.A. 2008. Defensins and the dynamic genome: what we can learn from structural variation at human chromosome band 8p23.1. Genome Res. 18: 1686-1697.
- Kohan, R. and Bower, C. 2008. Improving the health care experiences of families given the prenatal diagnosis of trisomy 18. J. Perinatol. 28: 719.
- 6. Shaw, J. 2008. Trisomy 18: a case study. Neonatal Netw. 27: 33-41.
- 7. Turleau, C. 2008. Monosomy 18p. Orphanet J. Rare Dis. 3: 4.
- 8. Witters, I. and Fryns, J.P. 2008. Trisomy 18 presenting with severe limb deformations. Prenat. Diagn. 28: 549-550.
- Edwards, S. and Waters, J.J. 2008. Prenatal diagnosis of monosomy 18p involving a jumping translocation. Prenat. Diagn. 28: 764-766.

CHROMOSOMAL LOCATION

Genetic locus: LOC284230 (human) mapping to 18p11.21.

SOURCE

LOC284230 (0-10) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of LOC284230 of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 100 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-84881 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

LOC284230 (Q-10) is recommended for detection of LOC284230 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for LOC284230 siRNA (h): sc-75454, LOC284230 shRNA Plasmid (h): sc-75454-SH and LOC284230 shRNA (h) Lentiviral Particles: sc-75454-V.

Molecular Weight of LOC284230: 12 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**