
SANTA CRUZ BIOTECHNOLOGY, INC.

RhoGEF p115/Lsc (C-19): sc-8492



BACKGROUND

The Ras superfamily of GTPases can be subdivided into the Ras, Rho/Rac, Sar, Rab, Arf and Ran subfamilies and controls multiple aspects of cell function, including cytoskeletal rearrangement, nuclear signaling and cell growth. The Ras superfamily of GTPases function as regulated switches that toggle between a biologically active GTP-bound and an inactive GDP-bound form. This activation is catalyzed by guanine nucleotide exchange factors (GEFs). The DbI-related proteins are a large family of structurally related molecules that have a common ability to catalyze GEF activity for specific members of the Ras family. DbI-related proteins include FGD1, RhoGEF p115/Lsc, Lfc, Lbc and Brx. RhoGEF p115/Lsc, Lbc and Lfc share sequence homology and show exchange activity toward Rho family GTPases. RhoGEF p115 (the human homolog of Lsc) catalyzes GEF activity for Rho but not Rac, Cdc42 or Ras GTPases.

CHROMOSOMAL LOCATION

Genetic locus: ARHGEF1 (human) mapping to 19q13.13; Arhgef1 (mouse) mapping to 7 A3.

SOURCE

RhoGEF p115/Lsc (C-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of RhoGEF p115 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-8492 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

RhoGEF p115/Lsc (C-19) is recommended for detection of RhoGEF p115 of human origin and Lsc of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RhoGEF p115 siRNA (h): sc-41734, Lsc siRNA (m): sc-41725, RhoGEF p115 shRNA Plasmid (h): sc-41734-SH, Lsc shRNA Plasmid (m): sc-41725-SH, RhoGEF p115 shRNA (h) Lentiviral Particles: sc-41734-V and Lsc shRNA (m) Lentiviral Particles: sc-41725-V.


Molecular Weight of RhoGEF p115/Lsc: 115 kDa.


Positive Controls: Jurkat whole cell lysate: sc-2204, RhoGEF p115 (h): 293T Lysate: sc-175428 or K-562 whole cell lysate: sc-2203.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunopre-cipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA





RhoGEF p115/Lsc (C-19): sc-8492. Western blot analysis of RhoGEF p115 expression in non-transfected 293T: sc-11752 (**A**), human RhoGEF p115 transfected 293T: sc-175428 (**B**), K-562 (**C**) and Jurkat (**D**) whole cell lysates. RhoGEF p115/Lsc (C-19): sc-8492. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

 Wang, Q., et al. 2004. Thrombin and lysophosphatidic acid receptors utilize distinct RhoGEFs in prostate cancer cells. J. Biol. Chem. 279: 28831-28834.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **RhoGEF p115 (C-9): sc-74565** or **Lsc (F-3): sc-374533**, our highly recommended monoclonal aternatives to RhoGEF p115/Lsc (C-19).