

ZNF407 (G-16): sc-85237



The Power to Question

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. As a member of the krueppel C₂H₂-type zinc-finger protein family, ZNF407 (zinc finger protein 407) is a 2,248 amino acid nuclear protein that contains 22 C₂H₂-type zinc fingers. The gene encoding ZNF407 maps to human chromosome 18, in a region that is frequently found to be affected in 18q deletion syndrome, a multiple-anomaly mental retardation syndrome. There are three isoforms of ZNF407 that are expressed as a result of alternative splicing events.

REFERENCES

1. Payre, F. and Vincent, A. 1988. Finger proteins and DNA-specific recognition: distinct patterns of conserved amino acids suggest different evolutionary modes. *FEBS Lett.* 234: 245-250.
2. Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
3. Rosenfeld, R. and Margalit, H. 1993. Zinc fingers: conserved properties that can distinguish between spurious and actual DNA-binding motifs. *J. Biomol. Struct. Dyn.* 11: 557-570.
4. Nuijten, I., Admiraal, R., Van Buggenhout, G., Cremers, C., Frijns, J.P., Smeets, D. and van Ravenswaaij-Arts, C. 2003. Congenital aural atresia in 18q deletion or de Grouchy syndrome. *Otol. Neurotol.* 24: 900-906.
5. Nusbaum, C., Zody, M.C., Borowsky, M.L., Kamal, M., Kodira, C.D., Taylor, T.D., Whittaker, C.A., Chang, J.L., Cuomo, C.A., Dewar, K., FitzGerald, M.G., Yang, X., Abouelleil, A., Allen, N.R., et al. 2005. DNA sequence and analysis of human chromosome 18. *Nature* 437: 551-555.
6. Dostal, A., Nemeckova, J., Gaillyova, R., Vranova, V., Zezulova, D., Lejska, M., Slapak, I., Dostalova, Z. and Kuglik, P. 2006. Identification of 2.3-Mb gene locus for congenital aural atresia in 18q22.3 deletion: a case report analyzed by comparative genomic hybridization. *Otol. Neurotol.* 27: 427-432.
7. Tariq, M., Chishti, M.S., Ali, G. and Ahmad, W. 2008. A novel locus for ectodermal dysplasia of hairs, nails and teeth type maps to chromosome 18q22.1-22.3. *Ann. Hum. Genet.* 72: 19-25.

CHROMOSOMAL LOCATION

Genetic locus: ZNF407 (human) mapping to 18q22.3; Zfp407 (mouse) mapping to 18 E4.

SOURCE

ZNF407 (G-16)-R is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the C-terminus of ZNF407 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-85237 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-85237 X, 100 µg/0.1 ml.

APPLICATIONS

ZNF407 (G-16)-R is recommended for detection of ZNF407 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ZNF family members.

ZNF407 (G-16) is also recommended for detection of ZNF407 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for ZNF407 siRNA (h): sc-76985, ZNF407 siRNA (m): sc-155710, ZNF407 shRNA Plasmid (h): sc-76985-SH, ZNF407 shRNA Plasmid (m): sc-155710-SH, ZNF407 shRNA (h) Lentiviral Particles: sc-76985-V and ZNF407 shRNA (m) Lentiviral Particles: sc-155710-V.

ZNF407 (G-16)-R X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of ZNF407: 247 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.