

BTBD3 (Y-15): sc-85311

BACKGROUND

BTBD3 (BTB/POZ domain-containing protein 3) is a 522 amino acid protein that contains one BTB/POZ domain. The BTB/POZ domain mediates homomeric and heteromeric POZ-POZ interactions and is common to transcriptional regulators involved in chromatin modeling. In several BTB/POZ containing proteins, including Bcl-6 and the promyelocytic leukemia zinc-finger (PLZF) oncoprotein, this domain interacts with the SMRT/N-CoR-mSin3A HDAC complex and is directly involved in repressing and silencing gene transcription. When this domain is deleted, as with the oncogenic PLZF-RAR chimera of promyelocytic leukemias, this transcriptional repression is attenuated. This suggests that BTBD3 may play a role in transcription regulation.

REFERENCES

1. Wong, C.W. and Privalsky, M.L. 1998. Components of the SMRT corepressor complex exhibit distinctive interactions with the POZ domain oncoproteins PLZF, PLZF-RAR α , and Bcl-6. *J. Biol. Chem.* 273: 27695-27702.
2. David, G., et al. 1998. Histone deacetylase associated with mSin3A mediates repression by the acute promyelocytic leukemia-associated PLZF protein. *Oncogene* 16: 2549-2556.
3. Huynh, K.D. and Bardwell, V.J. 1998. The Bcl-6 POZ domain and other POZ domains interact with the co-repressors N-CoR and SMRT. *Oncogene* 17: 2473-2484.
4. Ahmad, K.F., et al. 1998. Crystal structure of the BTB domain from PLZF. *Proc. Natl. Acad. Sci. USA* 95: 12123-12128.
5. Deltour, S., et al. 1999. Recruitment of SMRT/N-CoR-mSin3A-HDAC-repressing complexes is not a general mechanism for BTB/POZ transcriptional repressors: the case of HIC-1 and γ FDP-B. *Proc. Natl. Acad. Sci. USA* 96: 14831-14836.
6. Melnick, A., et al. 2002. Critical residues within the BTB domain of PLZF and Bcl-6 modulate interaction with corepressors. *Mol. Cell. Biol.* 22: 1804-1818.

CHROMOSOMAL LOCATION

Genetic locus: BTBD3 (human) mapping to 20p12.2; Btbd3 (mouse) mapping to 2 F3.

SOURCE

BTBD3 (Y-15) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of BTBD3 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-85311 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

BTBD3 (Y-15) is recommended for detection of BTBD3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other BTBD family members.

BTBD3 (Y-15) is also recommended for detection of BTBD3 in additional species, including canine, bovine, porcine and avian.

Suitable for use as control antibody for BTBD3 siRNA (h): sc-72667, BTBD3 siRNA (m): sc-141776, BTBD3 shRNA Plasmid (h): sc-72667-SH, BTBD3 shRNA Plasmid (m): sc-141776-SH, BTBD3 shRNA (h) Lentiviral Particles: sc-72667-V and BTBD3 shRNA (m) Lentiviral Particles: sc-141776-V.

Molecular Weight (predicted) of BTBD3: 58 kDa.


Molecular Weight (observed) of BTBD3: 41 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or mouse brain extract: sc-2253.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz MarkerTM compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz MarkerTM Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruzTM Mounting Medium: sc-24941.

DATA



BTBD3 (Y-15): sc-85311. Western blot analysis of BTBD3 expression in HeLa whole cell lysate (A) and mouse brain tissue extract (B).



BTBD3 (Y-15): sc-85311. Immunofluorescence staining of methanol-fixed HeLa cells showing membrane localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.