



LPLUNC3 (P-16): sc-85755

BACKGROUND

LPLUNC3 (long palate, lung and nasal epithelium carcinoma-associated protein 3), also known as C20orf185 or RYA3, is a 476 amino acid secreted protein that belongs to the BPI/LBP/PLUNC superfamily. Expressed in nasal septal epithelia, LPLUNC3 has the capacity to recognize and bind specific classes of odorants serving as a primary defense mechanism in the mucosa. LPLUNC3 is encoded by a gene which maps to a BPI/LBP/PLUNC gene cluster on human chromosome 20. Comprising approximately 2% of the human genome, chromosome 20 contains nearly 63 million bases that encode over 600 genes, some of which are associated with Creutzfeldt-Jakob disease, amyotrophic lateral sclerosis, spinal muscular atrophy, ring chromosome 20 epilepsy syndrome and Alagille syndrome.

REFERENCES

1. Bingle, C.D. and Craven, C.J. 2002. PLUNC: a novel family of candidate host defence proteins expressed in the upper airways and nasopharynx. *Hum. Mol. Genet.* 11: 937-943.
2. Zhang, B., Nie, X., Xiao, B., Xiang, J., Shen, S., Gong, J., Zhou, M., Zhu, S., Zhou, J., Qian, J., Lu, H., He, X., Li, X., Hu, G. and Li, G. 2003. Identification of tissue-specific genes in nasopharyngeal epithelial tissue and differentially expressed genes in nasopharyngeal carcinoma by suppression subtractive hybridization and cDNA microarray. *Genes Chromosomes Cancer* 38: 80-90.
3. Hou, J., Yashiro, K., Okazaki, Y., Saijoh, Y., Hayashizaki, Y. and Hamada, H. 2004. Identification of a novel left-right asymmetrically expressed gene in the mouse belonging to the BPI/PLUNC superfamily. *Dev. Dyn.* 229: 373-379.
4. LeClair, E.E., Nomellini, V., Bahena, M., Singleton, V., Bingle, L., Craven, C.J. and Bingle, C.D. 2004. Cloning and expression of a mouse member of the PLUNC protein family exclusively expressed in tongue epithelium. *Genomics* 83: 658-666.
5. Casado, B., Pannell, L.K., Iadarola, P. and Baraniuk, J.N. 2005. Identification of human nasal mucous proteins using proteomics. *Proteomics* 5: 2949-2959.
6. Flach, C.F., Qadri, F., Bhuiyan, T.R., Alam, N.H., Jennische, E., Lönnroth, I. and Holmgren, J. 2007. Broad upregulation of innate defense factors during acute cholera. *Infect. Immun.* 75: 2343-2350.
7. Vargas, P.A., Speight, P.M., Bingle, C.D., Barrett, A.W. and Bingle, L. 2008. Expression of PLUNC family members in benign and malignant salivary gland tumours. *Oral Dis.* 14: 613-619.

CHROMOSOMAL LOCATION

Genetic locus: C20orf185 (human) mapping to 20q11.21; RP23-244H7.9 (mouse) mapping to 2 H1.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

LPLUNC3 (P-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of LPLUNC3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-85755 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

LPLUNC3 (P-16) is recommended for detection of LPLUNC3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for LPLUNC3 siRNA (h): sc-75695, LPLUNC3 siRNA (m): sc-149024, LPLUNC3 shRNA Plasmid (h): sc-75695-SH, LPLUNC3 shRNA Plasmid (m): sc-149024-SH, LPLUNC3 shRNA (h) Lentiviral Particles: sc-75695-V and LPLUNC3 shRNA (m) Lentiviral Particles: sc-149024-V.

Molecular Weight of LPLUNC3: 50 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.