

## NIF-1 (N-20): sc-85792

### BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. NIF-1 (NRC-interacting factor 1), also known as NIF2 or ZNF335 (zinc finger protein 335), is a 1,342 amino acid nuclear protein belonging to the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc-finger protein family and exists as 2 alternatively spliced isoforms. NIF-1 is highly expressed in skeletal muscle, thymus, placenta and blood and contains thirteen C<sub>2</sub>H<sub>2</sub>-type zinc fingers. NIF-1 may regulate transcriptional activation through PRIP (peroxisome proliferator-activated receptor-interacting protein), a nuclear receptor co-activator that interacts with members of the steroid hormone and thyroid hormone/retinoid receptor subfamilies in a ligand-dependent or ligand-enhanced manner. NIF-1 is phosphorylated upon DNA damage by either ATM or ATR.

### REFERENCES

1. Kato, N., Shimotohno, K., VanLeeuwen, D. and Cohen, M. 1990. Human proviral mRNAs down regulated in choriocarcinoma encode a zinc finger protein related to Krüppel. *Mol. Cell. Biol.* 10: 4401-4405.
2. Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
3. Huebner, K., Druck, T., Croce, C.M. and Thiesen, H.J. 1991. Twenty-seven nonoverlapping zinc finger cDNAs from human T cells map to nine different chromosomes with apparent clustering. *Am. J. Hum. Genet.* 48: 726-740.
4. Zhu, Y., Kan, L., Qi, C., Kanwar, Y.S., Yeldandi, A.V., Rao, M.S. and Reddy, J.K. 2000. Isolation and characterization of peroxisome proliferator-activated receptor (PPAR) interacting protein (PRIP) as a co-activator for PPAR. *J. Biol. Chem.* 275: 13510-13516.
5. Mahajan, M.A., Murray, A. and Samuels, H.H. 2002. NRC-interacting factor 1 is a novel cotransducer that interacts with and regulates the activity of the nuclear hormone receptor co-activator NRC. *Mol. Cell. Biol.* 22: 6883-6894.
6. Mahajan, M.A., Murray, A., Levy, D. and Samuels, H.H. 2007. Nuclear receptor coregulator (NRC): mapping of the dimerization domain, activation of p53 and Stat2, and identification of the activation domain AD2 necessary for nuclear receptor signaling. *Mol. Endocrinol.* 21: 1822-1834.
7. Garapaty, S., Mahajan, M.A. and Samuels, H.H. 2008. Components of the CCR4-NOT complex function as nuclear hormone receptor co-activators via association with the NRC-interacting factor NIF-1. *J. Biol. Chem.* 283: 6806-6816.

### CHROMOSOMAL LOCATION

Genetic locus: ZNF335 (human) mapping to 20q13.12; Zfp335 (mouse) mapping to 2 H3.

### SOURCE

NIF-1 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of NIF-1 of human origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-85792 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-85792 X, 100 µg/0.1 ml.

### APPLICATIONS

NIF-1 (N-20) is recommended for detection of NIF-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NIF-1 (N-20) is also recommended for detection of NIF-1 in additional species, including equine, canine and porcine.

Suitable for use as control antibody for NIF-1 siRNA (h): sc-75914, NIF-1 siRNA (m): sc-149972, NIF-1 shRNA Plasmid (h): sc-75914-SH, NIF-1 shRNA Plasmid (m): sc-149972-SH, NIF-1 shRNA (h) Lentiviral Particles: sc-75914-V and NIF-1 shRNA (m) Lentiviral Particles: sc-149972-V.

NIF-1 (N-20) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of NIF-1: 145 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.