

PDRG (G-15): sc-85814

BACKGROUND

PDRG (p53 and DNA damage regulated), also known as PDRG1 or C20orf126, is a 133 amino acid protein that localizes to the cytoplasm and belongs to the prefoldin subunit β family. Expressed predominately in normal testicular tissue, PDRG is functionally induced by ultraviolet light and is thought to play a role in chaperone-mediated protein folding, possibly playing a role in cellular degradation. The gene encoding PDRG maps to human chromosome 20, which houses over 600 genes some of which are associated with Creutzfeldt-Jakob disease, amyotrophic lateral sclerosis, spinal muscular atrophy, ring chromosome 20 epilepsy syndrome and Alagille syndrome. Additionally, chromosome 20 contains a region with numerous genes which are thought important for seminal production and may be potential targets for male contraception.

REFERENCES

- Luo, X., Huang, Y. and Sheikh, M.S. 2003. Cloning and characterization of a novel gene PDRG that is differentially regulated by p53 and ultraviolet radiation. *Oncogene* 22: 7247-7257.
- Wong, K.K., Lu, K.H., Malpica, A., Bodurka, D.C., Shvartsman, H.S., Schmandt, R.E., Thornton, A.D., Deavers, M.T., Silva, E.G. and Gershenson, D.M. 2007. Significantly greater expression of ER, PR, and ECAD in advanced-stage low-grade ovarian serous carcinoma as revealed by immunohistochemical analysis. *Int. J. Gynecol. Pathol.* 26: 404-409.
- Horta, M.C., de Assis, L.A., de Souza, A.F., de Araújo, V.C., Gomez, R.S. and Aguiar, M.C. 2007. p53 and p21WAF1/CIP1 overexpression at the invasive front of lower lip squamous cell carcinoma. *J. Oral Pathol. Med.* 36: 88-92.
- Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610789. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Nam, E.J. and Kim, Y.T. 2008. Alteration of cell-cycle regulation in epithelial ovarian cancer. *Int. J. Gynecol. Cancer* 18: 1169-1182.
- Al-Joudi, F.S., Iskandar, Z.A. and Rusli, J. 2008. The expression of p53 in invasive ductal carcinoma of the breast: a study in the North-East States of Malaysia. *Med. J. Malaysia.* 63: 96-99.
- Wise-Draper, T.M., Morreale, R.J., Morris, T.A., Mintz-Cole, R.A., Hoskins, E.E., Balsitis, S.J., Husseinzadeh, N., Witte, D.P., Wikenheiser-Brokamp, K.A., Lambert, P.F. and Wells, S.I. 2009. DEK proto-oncogene expression interferes with the normal epithelial differentiation program. *Am. J. Pathol.* 174: 71-81.

CHROMOSOMAL LOCATION

Genetic locus: PDRG1 (human) mapping to 20q11.21; PdrG1 (mouse) mapping to 2 H1.

SOURCE

PDRG (G-15) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of PDRG of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-85814 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PDRG (G-15) is recommended for detection of PDRG of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PDRG (G-15) is also recommended for detection of PDRG in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for PDRG siRNA (h): sc-76099, PDRG siRNA (m): sc-152142, PDRG shRNA Plasmid (h): sc-76099-SH, PDRG shRNA Plasmid (m): sc-152142-SH, PDRG shRNA (h) Lentiviral Particles: sc-76099-V and PDRG shRNA (m) Lentiviral Particles: sc-152142-V.

Molecular Weight of PDRG: 15 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.