

S19BP (C-13): sc-86209

BACKGROUND

S19BP, also known as RPS19BP1 (ribosomal protein S19 binding protein 1) or AROS, is a 136 amino acid protein that localizes to the nucleolus and belongs to the AROS family. Expressed in a variety of tissues, S19BP interacts with ribosomal protein S19 and SIRT1 and is thought to directly regulate SIRT1 function, specifically enhancing the SIRT1-mediated deacetylation of p53 and inhibiting p53-induced transcriptional activity. The gene encoding S19BP maps to human chromosome 22, which houses over 500 genes and is the second smallest human chromosome. Mutations in several of the genes that map to chromosome 22 are involved in the development of Phelan-McDermid syndrome, neurofibromatosis type 2, autism and schizophrenia.

REFERENCES

- Gilbert, F. 1998. Disease genes and chromosomes: disease maps of the human genome. *Chromosome 22. Genet. Test.* 2: 89-97.
- Schwab, S.G. and Wildenauer, D.B. 1999. Chromosome 22 workshop report. *Am. J. Med. Genet.* 88: 276-278.
- Tsilchorozidou, T., Menko, F.H., Lalloo, F., Kidd, A., De Silva, R., Thomas, H., Smith, P., Malcolmson, A., Dore, J., Madan, K., Brown, A., Yovos, J.G., Tsaligopoulos, M., Vogiatzis, N., Baser, M.E., Wallace, A.J. and Evans, D.G. 2004. Constitutional rearrangements of chromosome 22 as a cause of neurofibromatosis 2. *J. Med. Genet.* 41: 529-534.
- Arinami, T. 2006. Analyses of the associations between the genes of 22q11 deletion syndrome and schizophrenia. *J. Hum. Genet.* 51: 1037-1045.
- Paylor, R., Glaser, B., Mupo, A., Ataliotis, P., Spencer, C., Sobotka, A., Sparks, C., Choi, C.H., Oghalai, J., Curran, S., Murphy, K.C., Monks, S., Williams, N., O'Donovan, M.C., Owen, M.J., Scambler, P.J. and Lindsay, E. 2006. Tbx1 haploinsufficiency is linked to behavioral disorders in mice and humans: implications for 22q11 deletion syndrome. *Proc. Natl. Acad. Sci. USA* 103: 7729-7734.
- Maeda, N., Toku, S., Kenmochi, N. and Tanaka, T. 2006. A novel nucleolar protein interacts with ribosomal protein S19. *Biochem. Biophys. Res. Commun.* 339: 41-46.
- Kim, E.J., Kho, J.H., Kang, M.R. and Um, S.J. 2007. Active regulator of SIRT1 cooperates with SIRT1 and facilitates suppression of p53 activity. *Mol. Cell* 28: 277-290.
- Hay, B.N. 2007. Deletion 22q11: spectrum of associated disorders. *Semin. Pediatr. Neurol.* 14: 136-139.
- Online Mendelian Inheritance in Man, OMIM[™]. 20007. Johns Hopkins University, Baltimore, MD. MIM Number: 610225. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: RPS19BP1 (human) mapping to 22q13.1; Rps19bp1 (mouse) mapping to 15 E1.

SOURCE

S19BP (C-13) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the C-terminus of Q86WX3 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-86209 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

S19BP (C-13) is recommended for detection of S19BP of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

S19BP (C-13) is also recommended for detection of S19BP in additional species, including equine.

Suitable for use as control antibody for S19BP siRNA (h): sc-76439, S19BP siRNA (m): sc-153192, S19BP shRNA Plasmid (h): sc-76439-SH, S19BP shRNA Plasmid (m): sc-153192-SH, S19BP shRNA (h) Lentiviral Particles: sc-76439-V and S19BP shRNA (m) Lentiviral Particles: sc-153192-V.

Molecular Weight of S19BP: 15 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.