

# DDT (T-14): sc-86407

## BACKGROUND

Macrophage migration inhibitory factor, known as MIF or glycosylation-inhibiting factor, is a secreted, homotrimeric, pro-inflammatory cytokine that modulates macrophage and T cell function and is an important regulator of host response to infection. MIF is expressed at sites of inflammation, which suggests that it plays a role in regulating macrophage function in host defense. The only known family member of MIF is D-dopachrome tautomerase (DDT), a protein that is thought to similarly play a role in the inflammation process. DDT is highly expressed in liver with lower levels in other organs, including heart, lung and pancreas. It resides in the cytoplasm as a homotrimer and converts 2-carboxy-2,3-dihydroindole-5, 6-quinone (D-dopachrome) into 5,6-dihydroxyindole. DDT requires the presence of an N-terminal proline residue for catalytic activity and is involved in the biosynthesis of melanin, an antioxidant. In response to liver damage, DDT has been shown to increase protein levels in order to accelerate melanin biosynthesis and protect the liver from oxidative stress.

## REFERENCES

- Weiser, W.Y., Temple, P.A., Witek-Giannotti, J.S., Remold, H.G., Clark, S.C. and David, J.R. 1989. Molecular cloning of a cDNA encoding a human macrophage migration inhibitory factor. *Proc. Natl. Acad. Sci. USA* 86: 7522-7526.
- Paralkar, V. and Wistow, G. 1994. Cloning the human gene for macrophage migration inhibitory factor (MIF). *Genomics* 19: 48-51.
- Bernhagen, J., Mitchell, R.A., Calandra, T., Voelter, W., Cerami, A. and Bucala, R. 1994. Purification, bioactivity, and secondary structure analysis of mouse and human macrophage migration inhibitory factor (MIF). *Biochemistry* 33: 14144-14155.
- Yoshida, H., Nishihira, J., Suzuki, M. and Hikichi, K. 1997. NMR characterization of physicochemical properties of rat D-dopachrome tautomerase. *Biochem. Mol. Biol. Int.* 42: 891-899.
- Nishihira, J., Fujinaga, M., Kuriyama, T., Suzuki, M., Sugimoto, H., Nakagawa, A., Tanaka, I. and Sakai, M. 1998. Molecular cloning of human D-dopachrome tautomerase cDNA: N-terminal proline is essential for enzyme activation. *Biochem. Biophys. Res. Commun.* 243: 538-544.
- Sugimoto, H., Taniguchi, M., Nakagawa, A., Tanaka, I., Suzuki, M. and Nishihira, J. 1999. Crystal structure of human D-dopachrome tautomerase, a homologue of macrophage migration inhibitory factor, at 1.54 Å resolution. *Biochemistry* 38: 3268-3279.
- Sonesson, B., Rosengren, E., Hansson, A.S. and Hansson, C. 2003. UVB-induced inflammation gives increased d-dopachrome tautomerase activity in blister fluid which correlates with macrophage migration inhibitory factor. *Exp. Dermatol.* 12: 278-282.

## CHROMOSOMAL LOCATION

Genetic locus: DDT (human) mapping to 22q11.23.

## SOURCE

DDT (T-14) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of DDT of human origin.

## PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-86407 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

DDT (T-14) is recommended for detection of DDT of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DDT siRNA (h): sc-77103, DDT shRNA Plasmid (h): sc-77103-SH and DDT shRNA (h) Lentiviral Particles: sc-77103-V.

Molecular Weight of DDT: 12 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **DDT (1G1): sc-517061**, our highly recommended monoclonal alternative to DDT (T-14).