



LDOC1L (N-13): sc-86516

BACKGROUND

LDOC1 (leucine zipper protein downregulated in cancer cells) is a 146 amino acid nuclear protein that contains a leucine zipper-like motif and a proline-rich region that shares marked similarity with an SH3-binding domain. The protein localizes to the nucleus and is downregulated in some cancer cell lines. It is thought to regulate the transcriptional response mediated by the nuclear factor κ B (NF κ B). The gene has been proposed as a tumor suppressor gene whose protein product may have an important role in the development and/or progression of some cancers. LDOC1L (leucine zipper protein downregulated in cancer cells-like), also known as MART6 (mammalian retrotransposon-derived protein 6), is a 239 amino acid protein that belongs to the LDOC1 family.

REFERENCES

- Nagasaki, K., Manabe, T., Hanzawa, H., Maass, N., Tsukada, T. and Yamaguchi, K. 1999. Identification of a novel gene, LDOC1, down-regulated in cancer cell lines. *Cancer Lett.* 140: 227-234.
- Nagasaki, K., Schem, C., von Kaisenberg, C., Biallek, M., Rösel, F., Jonat, W. and Maass, N. 2003. Leucine-zipper protein, LDOC1, inhibits NF κ B activation and sensitizes pancreatic cancer cells to apoptosis. *Int. J. Cancer* 105: 454-458.
- Chih, D.Y., Park, D.J., Gross, M., Idos, G., Vuong, P.T., Hirma, T., Chumakov, A.M., Said, J. and Koeffler, H.P. 2004. Protein partners of C/EBP ϵ . *Exp. Hematol.* 32: 1173-1181.
- Collins, J.E., Wright, C.L., Edwards, C.A., Davis, M.P., Grinham, J.A., Cole, C.G., Goward, M.E., Aguado, B., Mallya, M., Mokrab, Y., Huckle, E.J., Beare, D.M. and Dunham, I. 2004. A genome annotation-driven approach to cloning the human ORFeome. *Genome Biol.* 5: R84.
- Inoue, M., Takahashi, K., Niide, O., Shibata, M., Fukuzawa, M. and Ra, C. 2005. LDOC1, a novel MZF-1-interacting protein, induces apoptosis. *FEBS Lett.* 579: 604-608.
- Baffoe-Bonnie, A.B., Smith, J.R., Stephan, D.A., Schleutker, J., Carpten, J.D., Kainu, T., Gillanders, E.M., Matikainen, M., Teslovich, T.M., Tammela, T., Sood, R., Balshem, A.M., Scarborough, S.D., Xu, J., Isaacs, W.B., Trent, J.M., Kallioniemi, O.P. and Bailey-Wilson, J.E. 2005. A major locus for hereditary prostate cancer in Finland: localization by linkage disequilibrium of a haplotype in the HPCX region. *Hum. Genet.* 117: 307-316.
- Mizutani, K., Koike, D., Suetsugu, S. and Takenawa, T. 2005. WAVE3 functions as a negative regulator of LDOC1. *J. Biochem.* 138: 639-646.
- Ogawa, R., Ishiguro, H., Kuwabara, Y., Kimura, M., Mitsui, A., Mori, Y., Mori, R., Tomoda, K., Katada, T., Harada, K. and Fujii, Y. 2008. Identification of candidate genes involved in the radiosensitivity of esophageal cancer cells by microarray analysis. *Dis. Esophagus* 21: 288-297.

CHROMOSOMAL LOCATION

Genetic locus: LDOC1L (human) mapping to 22q13.31.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

LDOC1L (N-13) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping at the N-terminus of LDOC1L of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-86516 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

LDOC1L (N-13) is recommended for detection of LDOC1L of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for LDOC1L siRNA (h): sc-75415, LDOC1L shRNA Plasmid (h): sc-75415-SH and LDOC1L shRNA (h) Lentiviral Particles: sc-75415-V.

Molecular Weight of LDOC1L: 26 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.