

# SEZ6L (D-19): sc-86856

## BACKGROUND

SEZ6L (seizure related 6 homolog-like) is a 1,024 amino acid single-pass type I membrane protein that localizes to the endoplasmic reticulum. Widely expressed, SEZ6L is a candidate tumor suppressor gene and may contribute to specialized endoplasmic reticulum functions in neurons. SEZ6L contains three CUB domains and five sushi (CCP/SCR) domains, which may be involved in protein-protein interactions and signal transduction. The gene encoding SEZ6L is located on human chromosome 22, which houses over 500 genes and is the second smallest human chromosome. Mutations in several of the genes that map to chromosome 22 are involved in the development of Phelan-McDermid syndrome, neurofibromatosis type 2, autism and schizophrenia.

## REFERENCES

- Schwab, S.G. and Wildenauer, D.B. 1999. Chromosome 22 workshop report. *Am. J. Med. Genet.* 88: 276-278.
- Nishioka, M., Kohno, T., Takahashi, M., Niki, T., Yamada, T., Sone, S. and Yokota, J. 2000. Identification of a 428-kb homozygously deleted region disrupting the SEZ6L gene at 22q12.1 in a lung cancer cell line. *Oncogene* 19: 6251-6260
- Stefansson, B. and Brautigan, D.L. 2006. Protein phosphatase 6 subunit with conserved Sit4-associated protein domain targets I $\kappa$ B $\epsilon$ . *J. Biol. Chem.* 281: 22624-22634.
- Gorlov, I.P., Meyer, P., Liloglou, T., Myles, J., Boettger, M., Cassidy, A., Girard, L., Minna, J.D., ischer, R., Duffy, S., Spitz, M.R., Haeussinger, K., Kammerer, S., Cantor, C., Dierkesmann, R., Field, J.K. and Amos, C.I. 2007. Seizure 6-like (SEZ6L) gene and risk for lung cancer. *Cancer Res.* 67: 8406-8411.
- Yu, Z.L., Jiang, J.M., Wu, D.H., Xie, H.J., Jiang, J.J., Zhou, L., Peng, L. and Bao, G.S. 2007. Febrile seizures are associated with mutation of seizure-related (SEZ) 6, a brain-specific gene. *J. Neurosci. Res.* 85: 166-172.
- Hay, B.N. 2007. Deletion 22q11: spectrum of associated disorders. *Semin. Pediatr. Neurol.* 14: 136-139.
- Kang, G.H., Lee, S., Cho, N.Y., Gandamihardja, T., Long, T.I., Weisenberger, D.J., Campan, M. and Laird, P.W. 2008. DNA methylation profiles of gastric carcinoma characterized by quantitative DNA methylation analysis. *Lab. Invest.* 88: 161-170.

## CHROMOSOMAL LOCATION

Genetic locus: SEZ6L (human) mapping to 22q12.1.

## SOURCE

SEZ6L (D-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of SEZ6L of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-86856 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

SEZ6L (D-19) is recommended for detection of SEZ6L of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:5000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SEZ6L (D-19) is also recommended for detection of SEZ6L in additional species, including equine, canine and porcine.

Suitable for use as control antibody for SEZ6L siRNA (h): sc-76487, SEZ6L shRNA Plasmid (h): sc-76487-SH and SEZ6L shRNA (h) Lentiviral Particles: sc-76487-V.

Molecular Weight of SEZ6L: 112 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.