

TAF A5 (V-12): sc-86897

BACKGROUND

A group of small secreted proteins known as the TAF A family consists of five highly homologous genes: TAF A1, TAF A2, TAF A3, TAF A4 and TAF A5. Members of the TAF A family contain conserved cysteine residues at fixed positions and are highly expressed in brain. The TAF A family may be distantly related to a member of the CC-chemokine family known as MIP-1 α , and have been postulated to regulate nervous and immune cells of the brain as neurokines or chemokines. TAF A5 (chemokine-like protein TAF A-5), also known as FAM19A5 (family with sequence similarity 19 (chemokine (C-C motif)-like), member A5), is a 132 amino acid protein that belongs to the FAM19/TAF A family. A single-pass membrane protein, TAF A5 consists of three alternatively spliced variants that are encoded by a gene located on human chromosome 22. TAF A5 isoform 2 is a brain-specific secreted protein

REFERENCES

- Andersson, B., Wentland, M.A., Ricafrente, J.Y., Liu, W. and Gibbs, R.A. 1996. A "double adaptor" method for improved shotgun library construction. *Anal. Biochem.* 236: 107-113.
- Bonaldo, M.F., Lennon, G. and Soares, M.B. 1996. Normalization and subtraction: two approaches to facilitate gene discovery. *Genome Res.* 6: 791-806.
- Dunham, I., Shimizu, N., Roe, B.A., Chisoe, S., Hunt, A.R., Collins, J.E., Bruskiewich, R., Beare, D.M., Clamp, M., Smink, L.J., Ainscough, R., Almeida, J.P., Babbage, A., Bagguley, C., Bailey, J., Barlow, K., Bates, K.N., Beasley, O., Bird, C.P., Blakey, S., Bridgeman, A.M., Buck, D., Burgess, J., Burrill, W.D. and O'Brien, K.P. 1999. The DNA sequence of human chromosome 22. *Nature* 402: 489-495.
- Clark, H.F., Gurney, A.L., Abaya, E., Baker, K., Baldwin, D., Brush, J., Chen, J., Chow, B., Chui, C., Crowley, C., Currell, B., Deuel, B., Dowd, P., Eaton, D., Foster, J., Grimaldi, C., Gu, Q., Hass, P.E., Heldens, S., Huang, A., et al. 2003. The secreted protein discovery initiative (SPDI), a large-scale effort to identify novel human secreted and transmembrane proteins: a bioinformatics assessment. *Genome Res.* 13: 2265-2270.
- Tom Tang, Y., Emtage, P., Funk, W.D., Hu, T., Arterburn, M., Park, E.E., and Rupp, F. 2004. TAF A: a novel secreted family with conserved cysteine residues and restricted expression in the brain. *Genomics* 83: 727-734.
- Mick, E., Neale, B., Middleton, F.A., McGough, J.J. and Faraone, S.V. 2008. Genome-wide association study of response to methylphenidate in 187 children with attention-deficit/hyperactivity disorder. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 147B: 1412-1418.

CHROMOSOMAL LOCATION

Genetic locus: FAM19A5 (human) mapping to 22q13.32; Fam19a5 (mouse) mapping to 15 E3.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

TAF A5 (V-12) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of TAF A5 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-86897 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TAF A5 (V-12) is recommended for detection of TAF A5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with TAF A1 and TAF A4.

Suitable for use as control antibody for TAF A5 siRNA (h): sc-76629, TAF A5 siRNA (m): sc-154060, TAF A5 shRNA Plasmid (h): sc-76629-SH, TAF A5 shRNA Plasmid (m): sc-154060-SH, TAF A5 shRNA (h) Lentiviral Particles: sc-76629-V and TAF A5 shRNA (m) Lentiviral Particles: sc-154060-V.

Molecular Weight of TAF A5: 14 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.