

TLR4 (C-18): sc-8694

BACKGROUND

Six human homologs of the *Drosophila* Toll receptor were initially identified based on their sequence similarities and designated toll-like receptors (TLR). Toll receptors are involved in mediating dorsoventral polarization in the developing *Drosophila* embryo and also participate in the host immunity. The TLR family of proteins are characterized by a highly conserved Toll homology (TH) domain, which is essential for Toll-induced signal transduction. TLR1, as well as the other TLR family members, are type I transmembrane receptors that characteristically contain an extracellular domain consisting of several leucine-rich regions along with a single cytoplasmic Toll/IL-1R-like domain. TLR2 and TLR4 are activated in response to lipopolysaccharide (LPS) stimulation, which results in the activation and translocation of NF κ B and suggests that these receptors are involved in mediating inflammatory responses. Expression of TLR receptors is highest in peripheral blood leukocytes, macrophages, and monocytes. TLR6 is highly homologous to TLR1, sharing greater than 65% sequence identity, and, like other members of TLR family, it induces NF κ B signaling upon activation.

CHROMOSOMAL LOCATION

Genetic locus: TLR4 (human) mapping to 9q33.1.

SOURCE

TLR4 (C-18) is available as either goat (sc-8694) or rabbit (sc-8694-R) polyclonal affinity purified antibody raised against a peptide mapping near the C-terminus of TLR4 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-8694 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TLR4 (C-18) is recommended for detection of TLR4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TLR4 siRNA (h): sc-40260, TLR4 shRNA Plasmid (h): sc-40260-SH and TLR4 shRNA (h) Lentiviral Particles: sc-40260-V.

Molecular Weight of glycosylated TLR4: 95/120 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SELECT PRODUCT CITATIONS

- Ohnishi, T., et al. 2001. N-linked glycosylations at Asn 26 and Asn 114 of human MD-2 are required for toll-like receptor 4-mediated activation of NF κ B by lipopolysaccharide. *J. Immunol.* 167: 3354-3359.
- Farnell, M.B., et al. 2003. Oxidative burst mediated by toll-like receptors (TLR) and CD14 on avian heterophils stimulated with bacterial toll agonists. *Dev. Comp. Immunol.* 27: 423-429.
- Pivarcsi, A., et al. 2003. Expression and function of toll-like receptors 2 and 4 in human keratinocytes. *Int. Immunol.* 15: 721-730.
- Rounioja, S., et al. 2005. Mechanism of acute fetal cardiovascular depression after maternal inflammatory challenge in mouse. *Am. J. Pathol.* 166: 1585-1592.
- van Bruggen, R., et al. 2007. Complement receptor 3 and toll-like receptor 4 act sequentially in uptake and intracellular killing of unopsonized *Salmonella enterica serovar Typhimurium* by human neutrophils. *Infect. Immun.* 75: 2655-2660.
- Szebeni, B., et al. 2007. Increased mucosal expression of toll-like receptor TLR2 and TLR4 in coeliac disease. *J. Pediatr. Gastroenterol. Nutr.* 45: 187-193.
- Szczepanski, M., et al. 2007. Assessment of expression of toll-like receptors 2, 3 and 4 in laryngeal carcinoma. *Eur. Arch. Otorhinolaryngol.* 264: 525-530.
- Holmlund, U., et al. 2007. The novel inflammatory cytokine high mobility group box protein 1 (HMGB1) is expressed by human term placenta. *Immunology* 122: 430-437.
- Szebeni, B., et al. 2008. Increased expression of Toll-like receptor (TLR) 2 and TLR4 in the colonic mucosa of children with inflammatory bowel disease. *Clin. Exp. Immunol.* 151: 34-41.
- Aboussahoud, W., et al. 2010. Expression and function of Toll-like receptors in human endometrial epithelial cell lines. *J. Reprod. Immunol.* 84: 41-51.
- Liévin-Le Moal, V., et al. 2011. Apical expression of human full-length hCEACAM1-4L protein renders the madin darby canine kidney cells responsive to lipopolysaccharide leading to TLR4-dependent Erk1/2 and p38 MAPK signalling. *Cell. Microbiol.* 13: 764-785.
- Butti, E., et al. 2012. Subventricular zone neural progenitors protect striatal neurons from glutamatergic excitotoxicity. *Brain* 135: 3320-3335.
- Chan, S.T., et al. 2013. Quercetin enhances the antitumor activity of trichostatin A through upregulation of p53 protein expression *in vitro* and *in vivo*. *PLoS ONE* 8: e54255.



Try **TLR4 (25): sc-293072** or **TLR4 (76B357.1): sc-52962**, our highly recommended monoclonal alternatives to TLR4 (C-18). Also, for AC, HRP, FITC, PE, Alexa Fluor[®] 488 and Alexa Fluor[®] 647 conjugates, see **TLR4 (25): sc-293072**.