

Ataxin-1 (C-20): sc-8766

BACKGROUND

Ataxin-1, also designated spinocerebellar ataxia type 1 protein (Sca-1), is differentially expressed and localizes to both the cytoplasm and the nucleus. Mutations in Ataxin-1 are associated with the onset of the autosomal dominant neurodegenerative disorder spinocerebellar ataxia type 1 (SCA-1), which is characterized by progressive neuronal loss in the cerebellum, muscle wasting and ataxia. In Purkinje cells, where SCA-1 is predominantly observed, Ataxin-1 has been shown to directly associate with the Purkinje-enriched leucine-rich acidic nuclear protein (LANP) and the nuclear matrix-associated protein promyelocytic leukemia protein PML. In SCA-1, Ataxin-1 is mutated to encode a polyglutamine protein that forms nuclear aggregates, which interact significantly more strongly with LANP and contribute to the pathogenesis of SCA-1.

REFERENCES

1. Banfi, S., et al. 1994. Identification and characterization of the gene causing type 1 spinocerebellar ataxia. *Nat. Genet.* 7: 513-520.
2. Burchright, E.N., et al. 1995. SCA-1 transgenic mice: a model for neurodegeneration caused by an expanded CAG trinucleotide repeat. *Cell* 82: 937-948.
3. Burchright, E.N., et al. 1997. Identification of a self-association region within the SCA1 gene product, Ataxin-1. *Hum. Mol. Genet.* 6: 513-518.
4. Skinner, P.J., et al. 1997. Ataxin-1 with an expanded glutamine tract alters nuclear matrix-associated structures. *Nature* 389: 971-974.
5. Matilla, A., et al. 1997. The cerebellar leucine-rich acidic nuclear protein interacts with Ataxin-1. *Nature* 389: 974-978.
6. Klement, I.A., et al. 1998. Ataxin-1 nuclear localization and aggregation: role in polyglutamine-induced disease in SCA1 transgenic mice. *Cell* 95: 41-53.

CHROMOSOMAL LOCATION

Genetic locus: ATXN1 (human) mapping to 6p22.3; Atxn1 (mouse) mapping to 13 A5.

SOURCE

Ataxin-1 (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Ataxin-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-8766 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Ataxin-1 (C-20) is recommended for detection of Ataxin-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Ataxin-1 (C-20) is also recommended for detection of Ataxin-1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Ataxin-1 siRNA (h): sc-43624, Ataxin-1 siRNA (m): sc-40355, Ataxin-1 shRNA Plasmid (h): sc-43624-SH, Ataxin-1 shRNA Plasmid (m): sc-40355-SH, Ataxin-1 shRNA (h) Lentiviral Particles: sc-43624-V and Ataxin-1 shRNA (m) Lentiviral Particles: sc-40355-V.


Molecular Weight of Ataxin-1: 98 kDa.

Positive Controls: Ataxin-1 (m): 293T Lysate: sc-118599.

DATA



Ataxin-1 (C-20): sc-8766. Western blot analysis of Ataxin-1 expression in non-transfected: sc-117752 (A) and mouse Ataxin-1 transfected: sc-118599 (B) 293T whole cell lysates.



Ataxin-1 (C-20): sc-8766. Immunoperoxidase staining of formalin fixed, paraffin-embedded human fallopian tube tissue showing cytoplasmic staining of glandular cells.

SELECT PRODUCT CITATIONS

1. De Martino, I., et al. 2009. Regulation of microRNA expression by HMGA1 proteins. *Oncogene* 28: 1432-1442.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Ataxin-1 (E-4): sc-514953** or **Ataxin-1 (B-3): sc-365343**, our highly recommended monoclonal alternatives to Ataxin-1 (C-20).