

SCOT-t siRNA (h): sc-88311

BACKGROUND

SCOT-t (succinyl-CoA:3-ketoacid-coenzyme A transferase 2, mitochondrial), also known as OXCT2 (3-oxoacid CoA transferase 2), is a 517 amino acid protein that belongs to the 3-oxoacid CoA-transferase family. SCOT-t is a testis-specific succinyl-CoA:3-oxoacid CoA transferase, which catalyzes the reversible transfer of CoA from succinyl-CoA to acetoacetate in the first step of ketone body utilization. As a key enzyme for ketone body catabolism, SCOT-t transfers the CoA moiety from succinate to acetoacetate. Formation of the enzyme-CoA intermediate proceeds via an unstable anhydride species formed between the carboxylate groups of the enzyme and substrate. The SCOT-t protein has been detected in testis and spermatozoa, where it localized specifically to the mid-piece of sperm flagellum. The human SCOT-t protein shares 75.8% and 74.6% amino acid identity with mouse SCOT-t and human SCOT, respectively. The SCOT-t gene lacks an intron, and a nearby nonfunctional pseudogene has been identified in which nucleotides 745-762 and 778 are deleted. The entire SCOT-t transcription unit is located within an intron of the BMP-8 gene. The SCOT-t gene maps to human chromosome 1p34.2.

REFERENCES

- Lloyd, A.J. and Shoolingin-Jordan, P.M. 2001. Dimeric pig heart succinate-coenzyme A transferase uses only one subunit to support catalysis. *Biochemistry* 40: 2455-2467.
- Tanaka, H., Kohroki, J., Iguchi, N., Onishi, M. and Nishimune, Y. 2002. Cloning and characterization of a human orthologue of testis-specific succinyl CoA:3-oxoacid CoA transferase (SCOT-t) cDNA. *Mol. Hum. Reprod.* 8: 16-23.
- Cheung, J., Wilson, M.D., Zhang, J., Khaja, R., MacDonald, J.R., Heng, H.H., Koop, B.F. and Scherer, S.W. 2003. Recent segmental and gene duplications in the mouse genome. *Genome Biol.* 4: R47.
- Onishi, M., Yasunaga, T., Tanaka, H., Nishimune, Y. and Nozaki, M. 2004. Gene structure and evolution of testicular haploid germ cell-specific genes, *Oxct2a* and *Oxct2b*. *Genomics* 83: 647-657.
- Gregory, S.G., Barlow, K.F., McLay, K.E., Kaul, R., Swarbreck, D., Dunham, A., Scott, C.E., Howe, K.L., Woodfine, K., Spencer, C.C., Jones, M.C., Gillson, C., Searle, S., Zhou, Y., Kokocinski, F., McDonald, L., et al. 2006. The DNA sequence and biological annotation of human chromosome 1. *Nature* 441: 315-321.
- Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 610289. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Kuratomi, G., Iwamoto, K., Bundo, M., Kusumi, I., Kato, N., Iwata, N., Ozaki, N. and Kato, T. 2008. Aberrant DNA methylation associated with bipolar disorder identified from discordant monozygotic twins. *Mol. Psychiatry* 13: 429-441.
- Kumar, R.A., Sudi, J., Babatz, T.D., Brune, C.W., Oswald, D., Yen, M., Nowak, N.J., Cook, E.H., Christian, S.L. and Dobyns, W.B. 2010. A *de novo* 1p34.2 microdeletion identifies the synaptic vesicle gene *RIMS3* as a novel candidate for autism. *J. Med. Genet.* 47: 81-90.

CHROMOSOMAL LOCATION

Genetic locus: OXCT2 (human) mapping to 1p34.2.

PRODUCT

SCOT-t siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SCOT-t shRNA Plasmid (h): sc-88311-SH and SCOT-t shRNA (h) Lentiviral Particles: sc-88311-V as alternate gene silencing products.

For independent verification of SCOT-t (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-88311A, sc-88311B and sc-88311C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SCOT-t siRNA (h) is recommended for the inhibition of SCOT-t expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SCOT-t gene expression knockdown using RT-PCR Primer: SCOT-t (h)-PR: sc-88311-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.