

# MMP-11 (C-20): sc-8837

## BACKGROUND

The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, fibronectin, laminin and proteoglycan. Transcription of MMP genes is differentially activated by phorbol ester, lipopolysaccharide (LPS) or staphylococcal enterotoxin B (SEB). MMP catalysis requires both calcium and zinc. MMP-3, MMP-10 and MMP-11 (also designated stromelysin-1, -2 and -3) activate procollagenase. MMP-3 activation of procollagenase can occur via two pathways. Direct activation by MMP-3 is slow and activation by MMP-3 in conjunction with tissue or plasma proteinases is rapid. MMP-10 is expressed in small intestine, and it is expressed at lower levels in lung and heart. MMP-11 is specifically expressed in stromal cells of breast carcinomas and contributes to epithelial cell malignancies.

## REFERENCES

1. Saus, J., et al. 1988. The complete primary structure of human matrix metalloproteinase-3. Identity with stromelysin. *J. Biol. Chem.* 263: 6742-6745.
2. Suzuki, K., et al. 1990. Mechanisms of activation of tissue procollagenase by matrix metalloproteinase 3 (stromelysin). *Biochemistry* 29: 10261-10270.
3. Basset, P., et al. 1990. A novel metalloproteinase gene specifically expressed in stromal cells of breast carcinomas. *Nature* 348: 699-704.
4. Birkedal-Hansen, H., et al. 1993. Matrix metalloproteinases: a review. *Crit. Rev. Oral Biol. Med.* 4: 197-250.
5. Reinemer, P., et al. 1994. Structural implications for the role of the N terminus in the 'superactivation' of collagenases. A crystallographic study. *FEBS Lett.* 338: 227-233.
6. Knauper, V., et al. 1996. Activation of human neutrophil procollagenase by stromelysin 2. *Eur. J. Biochem.* 235: 187-191.
7. Machein, U., et al. 1997. Expression of several matrix metalloproteinase genes in human monocytic cells. *Adv. Exp. Med. Biol.* 421: 247-251.

## CHROMOSOMAL LOCATION

Genetic locus: MMP11 (human) mapping to 22q11.23; Mmp11 (mouse) mapping to 10 C1.

## SOURCE

MMP-11 (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of MMP-11 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-8837 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

MMP-11 (C-20) is recommended for detection of MMP-11 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MMP-11 (C-20) is also recommended for detection of MMP-11 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for MMP-11 siRNA (h): sc-35947, MMP-11 siRNA (m): sc-35948, MMP-11 shRNA Plasmid (h): sc-35947-SH, MMP-11 shRNA Plasmid (m): sc-35948-SH, MMP-11 shRNA (h) Lentiviral Particles: sc-35947-V and MMP-11 shRNA (m) Lentiviral Particles: sc-35948-V.

Molecular Weight of MMP-11: 60 kDa.

Positive Controls: ECV304 cell lysate: sc-2269, HeLa whole cell lysate: sc-2200 or HUV-EC-C whole cell lysate: sc-364180.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **MMP-11 (SL3.05): sc-58381**, our highly recommended monoclonal alternative to MMP-11 (C-20).