

AADACL3 siRNA (h): sc-88731

BACKGROUND

AADACL3 (arylacetylamide deacetylase-like 3), also known as RP11-474021.3, is a 350 amino acid protein that belongs to the "GDYG" lipolytic enzyme family and participates in hydrolase activity. Existing as two alternatively spliced isoforms, AADACL3 is encoded by a gene that maps to human chromosome 1p36.21. Chromosome 1, the largest human chromosome, makes up 8% of the human genome and contains about 260 million base pairs, which encode 3,000 genes. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

REFERENCES

- Blackwood, D.H., Fordyce, A., Walker, M.T., St Clair, D.M., Porteous, D.J. and Muir, W.J. 2001. Schizophrenia and affective disorders—cosegregation with a translocation at chromosome 1q42 that directly disrupts brain-expressed genes: clinical and P300 findings in a family. *Am. J. Hum. Genet.* 69: 428-433.
- Weise, A., Starke, H., Mrasek, K., Claussen, U. and Liehr, T. 2005. New insights into the evolution of chromosome 1. *Cytogenet. Genome Res.* 108: 217-222.
- Marzin, Y., Jamet, D., Douet-Guilbert, N., Morel, F., Le Bris, M.J., Morice, P., Abgrall, J.F., Berthou, C. and De Braekeleer, M. 2006. Chromosome 1 abnormalities in multiple myeloma. *Anticancer Res.* 26: 953-959.
- Gregory, S.G., Barlow, K.F., McLay, K.E., Kaul, R., Swarbreck, D., Dunham, A., Scott, C.E., Howe, K.L., Woodfine, K., Spencer, C.C., Jones, M.C., Gillson, C., Searle, S., Zhou, Y., et al. 2006. The DNA sequence and biological annotation of human chromosome 1. *Nature* 441: 315-321.
- Lindstrand, A., Malmgren, H., Sahlen, S., Xin, H., Schoumans, J. and Blennow, E. 2008. Molecular cytogenetic characterization of a constitutional, highly complex intrachromosomal rearrangement of chromosome 1, with 14 breakpoints and a 0.5 Mb submicroscopic deletion. *Am. J. Med. Genet. A* 146A: 3217-3222.
- Stacey, S.N., Gudbjartsson, D.F., Sulem, P., Bergthorsson, J.T., Kumar, R., Thorleifsson, G., Sigurdsson, A., Jakobsdottir, M., Sigurgeirsson, B., Benediksdottir, K.R., Thorisdottir, K., et al. 2008. Common variants on 1p36 and 1q42 are associated with cutaneous basal cell carcinoma but not with melanoma or pigmentation traits. *Nat. Genet.* 40: 1313-1318.
- Najfeld, V., Tripodi, J., Scalise, A., Silverman, L.R., Silver, R.T., Fruchtman, S. and Hoffman, R. 2010. Jumping translocations of the long arms of chromosome 1 in myeloid malignancies is associated with a high risk of transformation to acute myeloid leukaemia. *Br. J. Haematol.* 151: 288-921.
- SWISS-PROT/TrEMBL (Q5VUY0). World Wide Web URL: <http://www.uniprot.org/uniprot/Q5VUY0>

CHROMOSOMAL LOCATION

Genetic locus: AADACL3 (human) mapping to 1p36.21.

PRODUCT

AADACL3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AADACL3 shRNA Plasmid (h): sc-88731-SH and AADACL3 shRNA (h) Lentiviral Particles: sc-88731-V as alternate gene silencing products.

For independent verification of AADACL3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-88731A, sc-88731B and sc-88731C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AADACL3 siRNA (h) is recommended for the inhibition of AADACL3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AADACL3 gene expression knockdown using RT-PCR Primer: AADACL3 (h)-PR: sc-88731-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.