

# Tie-2 (H-176): sc-9026

## BACKGROUND

Receptor tyrosine kinases play key roles in signal transduction across cell surfaces in biological systems, including the vascular system. These receptors comprise a large and diverse family of catalytically related proteins that, on the basis of sequence and structural similarities, can be divided into several different evolutionary subfamilies. The cloning and characterization of Tie, a novel human endothelial cell surface receptor tyrosine kinase, has been reported. The extracellular domain of the predicted Tie protein product has an unusual multidomain structure consisting of a cluster of three epidermal growth factor homology motifs localized between two immunoglobulin-like loops, which are followed by three fibronectin type III repeats next to the transmembrane region. An additional member of this family has been identified as Tie-2 (also designated Tek). Tie-1 and Tie-2 have been shown to be encoded by distinct genes and to represent members of a new class of receptor tyrosine kinases.

## CHROMOSOMAL LOCATION

Genetic locus: TEK (human) mapping to 9p21.2; Tek (mouse) mapping to 4 C5.

## SOURCE

Tie-2 (H-176) is a rabbit polyclonal antibody raised against amino acids 25-200 mapping within the extracellular domain of Tie-2 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

Tie-2 (H-176) is recommended for detection of Tie-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Tie-2 (H-176) is also recommended for detection of Tie-2 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for Tie-2 siRNA (h): sc-36677, Tie-2 siRNA (m): sc-36678, Tie-2 shRNA Plasmid (h): sc-36677-SH, Tie-2 shRNA Plasmid (m): sc-36678-SH, Tie-2 shRNA (h) Lentiviral Particles: sc-36677-V and Tie-2 shRNA (m) Lentiviral Particles: sc-36678-V.

Molecular Weight of Tie-2: 140 kDa.

Positive Controls: HUV-EC-C whole cell lysate or ECV304 cell lysate: sc-2269.


## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## DATA



Tie-2 (H-176): sc-9026. Western blot analysis of Tie-2 expression in HUV-EC-C whole cell lysate.



Tie-2 (H-176): sc-9026. Western blot analysis of Tie-2 expression in ECV304 whole cell lysate.

## SELECT PRODUCT CITATIONS

- Nykanen, A.I., et al. 2003. Angiotensin-1 protects against the development of cardiac allograft arteriosclerosis. *Circulation* 107: 1308-1314.
- Nakayama, T., et al. 2004. Mast cell-derived angiotensin-1 plays a critical role in the growth of plasma cell tumors. *J. Clin. Invest.* 114: 1317-1325.
- Ramsden, J.D., et al. 2005. Complete inhibition of goiter in mice requires combined gene therapy modification of angiotensin, vascular endothelial growth factor, and fibroblast growth factor signaling. *Endocrinology* 146: 2895-2902.
- Glasker, S., et al. 2006. Hemangioblastomas share protein expression with embryonal hemangioblast progenitor cell. *Cancer Res.* 66: 4167-4172.
- Kim, K.L., et al. 2006. Interaction between Tie receptors modulates angiogenic activity of angiotensin2 in endothelial progenitor cells. *Cardiovasc. Res.* 72: 394-402.
- Patschan, D., et al. 2006. Dynamics of mobilization and homing of endothelial progenitor cells after acute renal ischemia: modulation by ischemic preconditioning. *Am. J. Physiol. Renal Physiol.* 291: F176-F185.
- Parborell, F., et al. 2008. Regulation of ovarian angiogenesis and apoptosis by GnRH-I analogs. *Mol. Reprod. Dev.* 75: 623-631.
- Kothmaier, H., et al. 2008. EGFR and PDGFR differentially promote growth in malignant epithelioid mesothelioma of short and long term survivors. *Thorax* 63: 345-351.
- Abramovich, D., et al. 2009. Spatiotemporal analysis of the protein expression of angiogenic factors and their related receptors during folliculogenesis in rats with and without hormonal treatment. *Reproduction* 137: 309-320.
- Scotti, L., et al. 2011. Administration of a gonadotropin-releasing hormone agonist affects corpus luteum vascular stability and development and induces luteal apoptosis in a rat model of ovarian hyperstimulation syndrome. *Mol. Cell. Endocrinol.* 335: 116-125.