

HSPA12A siRNA (h): sc-90428

BACKGROUND

The heat shock proteins (HSPs) comprise a group of highly conserved, abundantly expressed proteins with diverse functions, including the assembly and sequestering of multi-protein complexes, the transportation of nascent polypeptide chains across cellular membranes and the regulation of protein folding. HSPA12A (heat shock 70 kDa protein 12A) is a 675 amino acid protein that belongs to the HSP 70 family and is expressed in muscle, kidney and brain where it may play a role in cellular metabolism and regulation of neuronal function. The gene encoding HSPA12A maps to human chromosome 10, which houses over 1,200 genes and comprises nearly 4.5% of the human genome. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie-Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromic deafness, Wolman's syndrome, Cowden syndrome, multiple endocrine neoplasia type 2 and porphyria.

REFERENCES

1. Ishikawa, K., Nagase, T., Nakajima, D., Seki, N., Ohira, M., Miyajima, N., Tanaka, A., Kotani, H., Nomura, N. and Ohara, O. 1997. Prediction of the coding sequences of unidentified human genes. VIII. 78 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 4: 307-313.
2. Alimova-Kost, M.V., Imreh, S., Buchman, V.L. and Ninkina, N.N. 1998. Assignment1 of phosphotriesterase-related gene (PTER) to human chromosome band 10p12 by *in situ* hybridization. Cytogenet. Cell Genet. 83: 16-17.
3. Han, Z., Truong, Q.A., Park, S. and Breslow, J.L. 2003. Two HSP 70 family members expressed in atherosclerotic lesions. Proc. Natl. Acad. Sci. USA 100: 1256-1261.
4. Pongrac, J.L., Middleton, F.A., Peng, L., Lewis, D.A., Levitt, P. and Mirnics, K. 2004. Heat shock protein 12A shows reduced expression in the prefrontal cortex of subjects with schizophrenia. Biol. Psychiatry 56: 943-950.
5. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610701. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Brocchieri, L., Conway de Macario, E. and Macario, A.J. 2008. HSP 70 genes in the human genome: Conservation and differentiation patterns predict a wide array of overlapping and specialized functions. BMC Evol. Biol. 8: 19.

CHROMOSOMAL LOCATION

Genetic locus: HSPA12A (human) mapping to 10q25.3.

PRODUCT

HSPA12A siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HSPA12A shRNA Plasmid (h): sc-90428-SH and HSPA12A shRNA (h) Lentiviral Particles: sc-90428-V as alternate gene silencing products.

For independent verification of HSPA12A (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-90428A, sc-90428B and sc-90428C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HSPA12A siRNA (h) is recommended for the inhibition of HSPA12A expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HSPA12A gene expression knockdown using RT-PCR Primer: HSPA12A (h)-PR: sc-90428-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.