

CD28 (H-93): sc-9093

BACKGROUND

T cell proliferation and lymphokine production are triggered by occupation of the TCR by antigen, followed by a costimulatory signal that is delivered by a ligand expressed on antigen presenting cells. The B7-related cell surface proteins CD80 (B7-1) and CD86 (B7-2) are expressed on antigen presenting cells, bind the homologous T cell receptors CD28 and CTLA-4 (cytotoxic T lymphocyte-associated protein-4) and trigger costimulatory signals for optimal T cell activation. CTLA-4 shares 31% overall amino acid identity with CD28 and it has been proposed that CD28 and CTLA-4 are functionally redundant. SLAM is a novel receptor on T cells that, when engaged, potentiates T cell expansion in a CD28-independent manner. B7, also designated BB1, is another ligand or counterreceptor for CD28 and CTLA-4 that is expressed on the antigen-presenting cell.

REFERENCES

1. Freeman, G.J., et al. 1991. Structure, expression, and T cell costimulatory activity of the murine homologue of the human B lymphocyte activation antigen B7. *J. Exp. Med.* 174: 625-631.
2. Schwartz, R.H. 1992. Costimulation of T lymphocytes: the role of CD28, CTLA-4, and B7/BB1 in interleukin-2 production and immunotherapy. *Cell* 71: 1065-1068.

CHROMOSOMAL LOCATION

Genetic locus: CD28 (human) mapping to 2q33.2; Cd28 (mouse) mapping to 1 C2.

SOURCE

CD28 (H-93) is a rabbit polyclonal antibody raised against amino acids 19-111 of CD28 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CD28 (H-93) is recommended for detection of CD28 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CD28 siRNA (h): sc-29983, CD28 siRNA (m): sc-29982, CD28 shRNA Plasmid (h): sc-29983-SH, CD28 shRNA Plasmid (m): sc-29982-SH, CD28 shRNA (h) Lentiviral Particles: sc-29983-V and CD28 shRNA (m) Lentiviral Particles: sc-29982-V.

Molecular Weight of CD28 monomer: 44 kDa.


Molecular Weight of CD28 homodimer: 90 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CD28 (H-93): sc-9093. Western blot analysis of human recombinant CD28 fusion protein.

SELECT PRODUCT CITATIONS

1. Gorska, M.M., et al. 2004. UNC119, a novel activator of Lck/Fyn, is essential for T cell activation. *J. Exp. Med.* 199: 369-379.
2. Berg, M., et al. 2008. Regulation of CD28 expression on CD8⁺ T cells by CTLA-4. *J. Leukoc. Biol.* 83: 853-863.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **CD28 (CD28.2): sc-19655** or **CD28 (2Q1293): sc-70611**, our highly recommended monoclonal alternatives to CD28 (H-93).