

VCY siRNA (h): sc-91573

BACKGROUND

Human chromosome X contains nearly 153 million base pairs and houses over 1,000 genes, while human chromosome Y contains approximately 58 million base pairs and houses over 80 genes. The VCX and VCY (variable charge, X-linked and Y-linked, respectively) gene families are found on the X and Y chromosomes and encode small, highly charged proteins that are expressed specifically on male germ cells and may be involved in spermatogenesis. There are six members of the VCX/Y family, namely VCX-A, VCX-B, VCX-B1, VCX-C, VCY and VCY1B, all of which share a high degree of homology, with the exception of an amino acid sequence that is tandemly repeated in VCX proteins, but occurs only once in VCY proteins. Mutations or deletions in the genes encoding VCX proteins are associated with X-linked mental retardation. VCY (Variably charged protein Y), also known as testis-specific basic protein Y 1, is a 125 amino acid protein that is exclusively expressed in testis.

REFERENCES

1. Fukami, M., Kirsch, S., Schiller, S., Richter, A., Benes, V., Franco, B., Muroya, K., Rao, E., Merker, S., Niesler, B., Ballabio, A., Ansorge, W., Ogata, T. and Rappold, G.A. 2000. A member of a gene family on Xp22.3, VCX-A, is deleted in patients with X-linked nonspecific mental retardation. *Am. J. Hum. Genet.* 67: 563-573.
2. Lahn, B.T. and Page, D.C. 2000. A human sex-chromosomal gene family expressed in male germ cells and encoding variably charged proteins. *Hum. Mol. Genet.* 9: 311-319.
3. Zou, S.W., Zhang, J.C., Zhang, X.D., Miao, S.Y., Zong, S.D., Sheng, Q. and Wang, L.F. 2003. Expression and localization of VCX/Y proteins and their possible involvement in regulation of ribosome assembly during spermatogenesis. *Cell Res.* 13: 171-177.
4. Jiao, X., Wang, Z. and Kiledjian, M. 2006. Identification of an mRNA-decapping regulator implicated in X-linked mental retardation. *Mol. Cell.* 24: 713-722.
5. Cuevas-Covarrubias, S.A. and González-Huerta, L.M. 2008. Analysis of the VCX3A, VCX2 and VCX3B genes shows that VCX3A gene deletion is not sufficient to result in mental retardation in X-linked ichthyosis. *Br. J. Dermatol.* 158: 483-486.
6. Mochel, F., Missirian, C., Reynaud, R. and Moncla, A. 2008. Normal intelligence and social interactions in a male patient despite the deletion of NLGN4X and the VCX genes. *Eur. J. Med. Genet.* 51: 68-73.
7. Hansen, M.A., Nielsen, J.E., Retelska, D., Larsen, N. and Leffers, H. 2008. A shared promoter region suggests a common ancestor for the human VCX/Y, SPANX, and CSAG gene families and the murine CYPT family. *Mol. Reprod. Dev.* 75: 219-229.
8. Online Mendelian Inheritance in Man, OMIM[™]. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 400012. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: VCY (human) mapping to Yq11.221.

PRODUCT

VCY siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see VCY shRNA Plasmid (h): sc-91573-SH and VCY shRNA (h) Lentiviral Particles: sc-91573-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

VCY siRNA (h) is recommended for the inhibition of VCY expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor VCY gene expression knockdown using RT-PCR Primer: VCY (h)-PR: sc-91573-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.