

EphA7 (K-16): sc-917

BACKGROUND

The Eph subfamily represents the largest group of receptor protein tyrosine kinases identified to date. The Eph subfamily receptors of human origin (and their murine/avian homologs) include EphA1 (Eph), EphA2 (Eck), EphA3 (Hek4), EphA4 (Hek8), EphA5 (Hek7), EphA6 (Hek12), EphA7 (Hek11/MDK1), EphA8 (Hek3), EphB1 (Hek6), EphB2 (Hek5), EphB3 (Cek10, Hek2), EphB4 (Htk), EphB5 (Hek9) and EphB6 (Mep). EphAs are a family of receptor tyrosine kinases that are involved in axonal guidance during development. These receptors and their ligands, the ephrins, act via repulsive mechanisms to guide growing axons towards their appropriate targets and allow for the correct developmental connections to be made. Ligand binding to an Eph receptor results in tyrosine phosphorylation of the kinase domain, and repulsion of axonal growth cones and migrating cells. During neurulation, ephrin-A5 is coexpressed with its cognate receptor EphA7 in cells at the edges of the dorsal neural folds. Three different EphA7 splice variants, a full-length form and two truncated versions lacking kinase domains, are expressed in the neural folds.

REFERENCES

1. Ciossek, T., et al. 1995. Identification of alternatively spliced mRNAs encoding variants of MDK1, a novel receptor tyrosine kinase expressed in the murine nervous system. *Oncogene* 10: 97-108.
2. Fox, G.M., et al. 1995. DNA cloning and tissue distribution of five human Eph-like receptor protein-tyrosine kinases. *Oncogene* 10: 897-905.
3. Valenzuela, D.M., et al. 1995. Identification of full-length and truncated forms of Ehk-3, a novel member of the Eph receptor tyrosine kinase family. *Oncogene* 10: 1573-1580.
4. Holmberg, J., et al. 2000. Regulation of repulsion versus adhesion by different splice forms of an Eph receptor. *Nature* 408: 203-206.

CHROMOSOMAL LOCATION

Genetic locus: EPHA7 (human) mapping to 6q16.1; EphA7 (mouse) mapping to 4 A4.

SOURCE

EphA7 (K-16) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping at the N-terminus of EphA7 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-917 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

EphA7 (K-16) is recommended for detection of EphA7 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

EphA7 (K-16) is also recommended for detection of EphA7 in additional species, including equine, canine, bovine, porcine and avian.


Suitable for use as control antibody for EphA7 siRNA (h): sc-39941, EphA7 siRNA (m): sc-39942, EphA7 shRNA Plasmid (h): sc-39941-SH, EphA7 shRNA Plasmid (m): sc-39942-SH, EphA7 shRNA (h) Lentiviral Particles: sc-39941-V and EphA7 shRNA (m) Lentiviral Particles: sc-39942-V.

Molecular Weight (predicted) of EphA7: 112 kDa.


Molecular Weight (observed) of EphA7: 93 kDa.

Positive Controls: EphA7 (h2): 293T Lysate: sc-158479, A-10 cell lysate: sc-3806 or HeLa whole cell lysate: sc-2200.

DATA



EphA7 (K-16): sc-917. Western blot analysis of EphA7 expression in non-transfected: sc-117752 (A) and truncated human EphA7 transfected: sc-158479 (B) 293T whole cell lysates.



EphA7 (K-16): sc-917. Western blot analysis of EphA7 expression in rat small intestine tissue extract.

SELECT PRODUCT CITATIONS

1. Halford, M.M., et al. 2000. Ryk-deficient mice exhibit craniofacial defects associated with perturbed Eph receptor crosstalk. *Nat. Genet.* 25: 414-418.
2. Hu, Z., et al. 2003. Corpus callosum deficiency in transgenic mice expressing a truncated ephrin-A receptor. *J. Neurosci.* 23: 10963-10970.
3. Iwasa, H., et al. 2003. Mitogen-activated protein kinase p38 defines the common senescence-signalling pathway. *Genes Cells* 8: 131-144.
4. Dawson, D.W., et al. 2007. Global DNA methylation profiling reveals silencing of a secreted form of EphA7 in mouse and human germinal center B-cell lymphomas. *Oncogene* 26: 4243-4252.

MONOS
Satisfaction
Guaranteed

Try **EphA7 (E-7): sc-393973** or **EphA7 (F-10): sc-393974**, our highly recommended monoclonal alternatives to EphA7 (K-16).