

# SKAP55 (C-17): sc-9176

## BACKGROUND

Fyb (Fyn binding protein) and the anchoring proteins SKAP55 and SKAP55-R (SKAP55-related protein) associate with the tyrosine kinase p59fyn. SKAP55 and SKAP55-R bind to Fyb through their SH3 domains and function as substrates for p59Fyn in resting T cells. SKAP55 contains an N-terminal pleckstrin homology domain and a C-terminal SH3 domain binding motif of adjacent arginine and lysine residues followed by tandem tyrosines (i.e. RKxxYxxY). SKAP55-R, similar in overall structure to SKAP55, contains a coiled-coil N-terminal domain. SKAP55 associates with SLAP-130, another component of the Fyn complex, which plays a role in the regulation of signaling events initiated by lymphocyte antigen receptors leading up to T cell activation. The human SKAP55 gene maps to chromosome 17q21.32 and encodes a 359 amino acid protein.

## CHROMOSOMAL LOCATION

Genetic locus: SCAP1 (human) mapping to 17q21.32.

## SOURCE

SKAP55 (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of SKAP55 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9176 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

SKAP55 (C-17) is recommended for detection of SKAP55 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SKAP55 (C-17) is also recommended for detection of SKAP55 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for SKAP55 siRNA (h): sc-40599, SKAP55 shRNA Plasmid (h): sc-40599-SH and SKAP55 shRNA (h) Lentiviral Particles: sc-40599-V.


Molecular Weight of SKAP55: 55 kDa.

Positive Controls: SKAP55 (h): 293T Lysate: sc-116192, Jurkat whole cell lysate: sc-2204 or SUP-T1 whole cell lysate: sc-364796.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



SKAP55 (C-17): sc-9176. Western blot analysis of SKAP55 expression in non-transfected: sc-117752 (A) and human SKAP55 transfected: sc-116192 (B) 293T whole cell lysates.



SKAP55 (C-17): sc-9176. Western blot analysis of SKAP55 expression in non-transfected: sc-117752 (A) and human SKAP55 transfected: sc-116192 (B) 293T whole cell lysates.

## SELECT PRODUCT CITATIONS

- Bourette, R.P., et al. 2005. Macrophage colony-stimulating factor receptor induces tyrosine phosphorylation of SKAP55-R adaptor and its association with Actin. *Cell. Signal.* 17: 941-949.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **SKAP55 (E-8): sc-390458** or **SKAP55 (35): sc-136068**, our highly recommended monoclonal alternatives to SKAP55 (C-17).