

ZDHHC11 siRNA (h): sc-91779

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. ZDHHC11 (zinc finger, DHHC domain containing 11), also known as ZNF399, is a 412 amino acid multi-pass membrane protein that contains one DHHC-type zinc finger and is thought to function as a palmitoyl-transferase, catalyzing the transformation of palmitoyl-CoA and a cysteine-conjugated protein to a S-palmitoyl protein and free CoA. ZDHHC11 may be a potential biomarker identifying high-risk patients with disease progression in bladder cancer. The gene encoding ZDHHC11 maps to human chromosome 5, which contains 181 million base pairs and comprises nearly 6% of the human genome.

REFERENCES

1. Putilina, T., et al. 1999. The DHHC domain: a new highly conserved cysteine-rich motif. *Mol. Cell. Biochem.* 195: 219-226.
2. Roth, A.F., et al. 2002. The yeast DHHC cysteine-rich domain protein Akr1p is a palmitoyl transferase. *J. Cell Biol.* 159: 23-28.
3. Ohno, Y., et al. 2006. Intracellular localization and tissue-specific distribution of human and yeast DHHC cysteine-rich domain-containing proteins. *Biochim. Biophys. Acta* 1761: 474-483.
4. Mitchell, D.A., et al. 2006. Protein palmitoylation by a family of DHHC protein S-acyltransferases. *J. Lipid Res.* 47: 1118-1127.
5. Yamamoto, Y., et al. 2007. Gain of 5p15.33 is associated with progression of bladder cancer. *Oncology* 72: 132-138.

CHROMOSOMAL LOCATION

Genetic locus: ZDHHC11 (human) mapping to 5p15.33.

PRODUCT

ZDHHC11 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZDHHC11 shRNA Plasmid (h): sc-91779-SH and ZDHHC11 shRNA (h) Lentiviral Particles: sc-91779-V as alternate gene silencing products.

For independent verification of ZDHHC11 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-91779A and sc-91779B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZDHHC11 siRNA (h) is recommended for the inhibition of ZDHHC11 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZDHHC11 gene expression knockdown using RT-PCR Primer: ZDHHC11 (h)-PR: sc-91779-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.