

# BRMS1L siRNA (h): sc-92226

## BACKGROUND

BRMS1L (breast cancer metastasis-suppressor 1-like) is a 323 amino acid protein that localizes to the nucleus and exists as a component of the mSin3A/HDAC1 (histone deacetylase) complex. Sharing similarity with BRMS1, BRMS1L is involved in HDAC1-dependent transcriptional repression and, in lung cancer tissue, functions to inhibit cell growth, suggesting a role in tumor suppression. The gene encoding BRMS1L maps to human chromosome 14, which houses over 700 genes and comprises nearly 3.5% of the human genome. Chromosome 14 encodes the presenilin 1 (PSEN1) gene, which is one of the three key genes associated with the development of Alzheimer's disease (AD). The SERPINA1 gene is also located on chromosome 14 and, when defective, leads to the genetic disorder  $\alpha$ 1-antitrypsin deficiency, which is characterized by severe lung complications and liver dysfunction.

## REFERENCES

1. Kuzmichev, A., Zhang, Y., Erdjument-Bromage, H., Tempst, P. and Reinberg, D. 2002. Role of the Sin3-histone deacetylase complex in growth regulation by the candidate tumor suppressor p33<sup>ING1</sup>. *Mol. Cell. Biol.* 22: 835-848.
2. Nikolaev, A.Y., Papanikolaou, N.A., Li, M., Qin, J. and Gu, W. 2004. Identification of a novel BRMS1-homologue protein p40 as a component of the mSin3A/p33<sup>ING1b</sup>/HDAC1 deacetylase complex. *Biochem. Biophys. Res. Commun.* 323: 1216-1222.
3. Meehan, W.J., Samant, R.S., Hopper, J.E., Carrozza, M.J., Shevde, L.A., Workman, J.L., Eckert, K.A., Verderame, M.F. and Welch, D.R. 2004. Breast cancer metastasis suppressor 1 (BRMS1) forms complexes with retinoblastoma-binding protein 1 (RBP1) and the mSin3 histone deacetylase complex and represses transcription. *J. Biol. Chem.* 279: 1562-1569.
4. Albani, D., Roiter, I., Artuso, V., Batelli, S., Prato, F., Pesaresi, M., Galimberti, D., Scarpini, E., Bruni, A., Franceschi, M., Piras, M.R., Confalonni, A. and Forloni, G. 2007. Presenilin-1 mutation E318G and familial Alzheimer's disease in the Italian population. *Neurobiol. Aging* 28: 1682-1688.
5. Cruz, P.E., Mueller, C. and Flotte, T.R. 2007. The promise of gene therapy for the treatment of  $\alpha$ -1 antitrypsin deficiency. *Pharmacogenomics* 8: 1191-1198.

## CHROMOSOMAL LOCATION

Genetic locus: BRMS1L (human) mapping to 14q13.2.

## PRODUCT

BRMS1L siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see BRMS1L shRNA Plasmid (h): sc-92226-SH and BRMS1L shRNA (h) Lentiviral Particles: sc-92226-V as alternate gene silencing products.

For independent verification of BRMS1L (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-92226A, sc-92226B and sc-92226C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

BRMS1L siRNA (h) is recommended for the inhibition of BRMS1L expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

BRMS1L (E-2): sc-518200 is recommended as a control antibody for monitoring of BRMS1L gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor BRMS1L gene expression knockdown using RT-PCR Primer: BRMS1L (h)-PR: sc-92226-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.