

PPIL5 siRNA (h): sc-92289

BACKGROUND

Cyclophilins are conserved, ubiquitous and abundant cytosolic peptidylprolyl cis-trans isomerases that accelerate the isomerization of XaaPro peptide bonds and the refolding of proteins. PPIL5 (peptidylprolyl isomerase (cyclophilin)-like 5), also known as LRR-repeat protein 1 (LRR-1) or 4-1BB-mediated-signaling molecule (4-1BB_{lr}), is a 414 amino acid protein involved in protein modification and ubiquitination. PPIL5 has been found to negatively regulate CD137-mediated signaling cascades by interacting with the cytoplasmic domain of CD137 (also known as 4-1BB), which leads to activation of JNK1 and NFκB. While ubiquitously expressed, highest expression of PPIL5 is found in skeletal muscle and heart. PPIL5 contains five LRR (leucine-rich) repeats, and exists as two alternatively spliced isoforms. The gene encoding PPIL5 maps to human chromosome 14q21.3.

REFERENCES

1. Kwon, B., Moon, C.H., Kang, S., Seo, S.K. and Kwon, B.S. 2000. 4-1BB: still in the midst of darkness. *Mol. Cells* 10: 119-126.
2. Jang, L.K., Lee, Z.H., Kim, H.H., Hill, J.M., Kim, J.D. and Kwon, B.S. 2001. A novel leucine-rich repeat protein (LRR-1): potential involvement in 4-1BB-mediated signal transduction. *Mol. Cells* 12: 304-312.
3. Kamura, T., Maenaka, K., Kotoshiba, S., Matsumoto, M., Kohda, D., Conaway, R.C., Conaway, J.W. and Nakayama, K.I. 2004. VHL-box and SOCS-box domains determine binding specificity for CUL2-Rbx1 and CUL5-Rbx2 modules of ubiquitin ligases. *Genes Dev.* 18: 3055-3065.
4. Nebert, D.W., Sophos, N.A., Vasiliou, V. and Nelson, D.R. 2004. Cyclophilin nomenclature problems, or, "a visit from the sequence police". *Hum. Genomics* 1: 381-388.
5. Valle, C., Troiani, A.R., Lazzaretti, P., Bouvier, J., Cioli, D. and Klinkert, M.O. 2005. Molecular and biochemical characterization of a protein cyclophilin from the nematode *Haemonchus contortus* (P). *Parasitol. Res.* 96: 199-205.
6. Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 609193. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: LRR1 (human) mapping to 14q21.3.

PRODUCT

PPIL5 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PPIL5 shRNA Plasmid (h): sc-92289-SH and PPIL5 shRNA (h) Lentiviral Particles: sc-92289-V as alternate gene silencing products.

For independent verification of PPIL5 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-92289A, sc-92289B and sc-92289C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PPIL5 siRNA (h) is recommended for the inhibition of PPIL5 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PPIL5 gene expression knockdown using RT-PCR Primer: PPIL5 (h)-PR: sc-92289-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.