

DAF-2 (cN-17): sc-9231

BACKGROUND

Several proteins involved in regulating the aging process in *C. elegans* have been identified. DAF-2, DAF-16 and AGE-1 (also known as DAF-23) regulate lifespan via an Insulin-signaling pathway. In specific, decreases in DAF-2 signaling induce metabolic and developmental changes, as in mammalian metabolic control by the Insulin receptor. DAF-16 encodes a member of the hepatocyte nuclear factor 3 (HNF-3)/forkhead family of transcriptional regulators. In humans HNF-3 activity is antagonized by Insulin, causing the down regulation of developmental genes, raising the possibility that aspects of the DAF-16 regulatory system have been conserved. The gene AGE-1 encodes a homologue of mammalian phosphatidylinositol-3-OH kinase (PI(3)K) catalytic subunits and is required for non-dauer development and normal senescence. CLK-1, a homolog of the yeast COQ7/CAT5 protein, is thought to exert its effects on longevity via the synthesis of ubiquinone, an essential component of electron transport.

REFERENCES

- Wilson, R., Ainscough, R., Anderson, K., Baynes, C., Berks, M., Bonfield, J., Burton, J., Connell, M., Copsey, T., Cooper, J., et al. 1994. 2.2 Mb of contiguous nucleotide sequence from chromosome III of *C. elegans*. *Nature* 368: 32-38.
- Morris, J.Z., Tissenbaum, H.A. and Ruvkun, G. 1996. A phosphatidylinositol-3-OH kinase family member regulating longevity and diapause in *Caenorhabditis elegans*. *Nature* 382: 536-539.
- Ewbank, J.J., Barnes, T.M., Lakowski, B., Lussier, M., Bussey, H. and Hekimi, S. 1997. Structural and functional conservation of the *Caenorhabditis elegans* timing gene CLK-1. *Science* 275: 980-983.
- Kimura, K.D., Tissenbaum, H.A., Liu, Y. and Ruvkun, G. 1997. DAF-2, an Insulin receptor-like gene that regulates longevity and diapause in *Caenorhabditis elegans*. *Science* 277: 942-946.
- Lin, K., Dorman, J.B., Rodan, A. and Kenyon, C. 1997. DAF-16: An HNF-3/forkhead family member that can function to double the life-span of *Caenorhabditis elegans*. *Science* 278:1319-1322.
- Vajo, Z., King, L.M., Jonassen, T., Wilkin, D.J., Ho, N., Munnich, A., Clarke, C.F. and Francomano, C.A. 1999. Conservation of the *Caenorhabditis elegans* timing gene CLK-1 from yeast to human: a gene required for ubiquinone biosynthesis with potential implications for aging. *Mamm. Genome* 10: 1000-1004.
- Moeslein, F.M., Myers, M.P. and Landreth, G.E. 1999. The CLK family kinases, CLK-1 and CLK-2, phosphorylate and activate the tyrosine phosphatase, PTP-1B. *J. Biol. Chem.* 274: 26697-26704.

SOURCE

DAF-2 (cN-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of DAF-2 of *C. elegans* origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9231 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

DAF-2 (cN-17) is recommended for detection of DAF-2 of *Caenorhabditis elegans* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.