

TSCOT siRNA (h): sc-92745

BACKGROUND

TSCOT (thymic stromal cotransporter homolog) is a 475 amino acid multi-pass membrane protein that belongs to the SLC46A family. TSCOT is believed to act as a cell membrane transporter and is likely to be found glycosylated. TSCOT is encoded by a gene that maps to human chromosome 9q32, which consists of about 145 million bases and 4% of the human genome and encodes nearly 900 genes. Considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster.

REFERENCES

- Chen, C., Kim, M.G., Soo Lyu, M., Kozak, C.A., Schwartz, R.H. and Flomerfelt, F.A. 2000. Characterization of the mouse gene, human promoter and human cDNA of TSCOT reveals strong interspecies homology. *Biochim. Biophys. Acta* 1493: 159-169.
- Kim, M.G., Flomerfelt, F.A., Lee, K.N., Chen, C. and Schwartz, R.H. 2000. A putative 12 transmembrane domain cotransporter expressed in thymic cortical epithelial cells. *J. Immunol.* 164: 3185-3192.
- Obermann, H., Wingbermühle, A., Münz, S. and Kirchhoff, C. 2003. A putative 12-transmembrane domain cotransporter associated with apical membranes of the epididymal duct. *J. Androl.* 24: 542-556.
- Yang, S.J., Ahn, S., Park, C.S., Choi, S. and Kim, M.G. 2005. Identifying subpopulations of thymic epithelial cells by flow cytometry using a new specific thymic epithelial marker, Ly110. *J. Immunol. Methods* 297: 265-270.
- Engelmark, M.T., Ivansson, E.L., Magnusson, J.J., Gustavsson, I.M., Beskow, A.H., Magnusson, P.K. and Gyllensten, U.B. 2006. Identification of susceptibility loci for cervical carcinoma by genome scan of affected sib-pairs. *Hum. Mol. Genet.* 15: 3351-3360.
- Engelmark, M.T., Ivansson, E.L., Magnusson, J.J., Gustavsson, I.M., Wyöni, P.I., Ingman, M., Magnusson, P.K. and Gyllensten, U.B. 2008. Polymorphisms in 9q32 and TSCOT are linked to cervical cancer in affected sib-pairs with high mean age at diagnosis. *Hum. Genet.* 123: 437-443.
- Ahn, S., Lee, G., Yang, S.J., Lee, D., Lee, S., Shin, H.S., Kim, M.C., Lee, K.N., Palmer, D.C., Theoret, M.R., Jenkinson, E.J., Anderson, G., Restifo, N.P. and Kim, M.G. 2008. TSCOT⁺ thymic epithelial cell-mediated sensitive CD4 tolerance by direct presentation. *PLoS Biol.* 6: e191.
- Bateup, H.S., Takasaki, K.T., Saulnier, J.L., Deneffrio, C.L. and Sabatini, B.L. 2011. Loss of Tsc1 *in vivo* impairs hippocampal mGluR-LTD and increases excitatory synaptic function. *J. Neurosci.* 31: 8862-8869.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: SLC46A2 (human) mapping to 9q32.

PRODUCT

TSCOT siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TSCOT shRNA Plasmid (h): sc-92745-SH and TSCOT shRNA (h) Lentiviral Particles: sc-92745-V as alternate gene silencing products.

For independent verification of TSCOT (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-92745A, sc-92745B and sc-92745C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TSCOT siRNA (h) is recommended for the inhibition of TSCOT expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TSCOT gene expression knockdown using RT-PCR Primer: TSCOT (h)-PR: sc-92745-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.