

DAF-3 (cC-19): sc-9290

BACKGROUND

A variety of growth factor signaling molecules have been shown to regulate *C. elegans* development, including members of the EGF, FGF and TGF β superfamilies. These factors bind to specific receptors and transduce extracellular signals to the nucleus. Receptor tyrosine kinase/Ras pathways also play a critical role in cell signaling and are responsible for proper vulval development. The DAF family proteins play a role in the control of dauer larva formation, a developmentally arrested, non-feeding dispersal stage normally formed in response to overcrowding and limited food. DAF-1, DAF-3, DAF-4 and DAF-7 are members of the TGF β superfamily. DAF-18 is the *C. elegans* homologue of PTEN and acts as a component of the Insulin-like signalling pathway.

REFERENCES

- Georgi, L.L., Albert, P.S. and Riddle, D.L. 1990. DAF-1, a *C. elegans* gene controlling dauer larva development, encodes a novel receptor protein kinase. *Cell* 61: 635-645.
- Carpenter, G. 1993. EGF: new tricks for an old growth factor. *Curr. Opin. Cell Biol.* 5: 261-264.
- Sternberg, P.W., Lesa, G., Lee, J., Katz, W.S., Yoon, C., Clandinin, T.R., Huang, L.S., Chamberlin, H.M. and Jongeward, G. 1995. LET-23-mediated signal transduction during *Caenorhabditis elegans* development. *Mol. Reprod. Dev.* 42: 523-528.
- Kayne, P.S. and Sternberg, P.W. 1995. Ras pathways in *Caenorhabditis elegans*. *Curr. Opin. Genet. Dev.* 5: 38-43.
- Ren, P., Lim, C.S., Johnsen, R., Albert, P.S., Pilgrim, D. and Riddle, D.L. 1996. Control of *C. elegans* larval development by neuronal expression of a TGF β homologue. *Science* 274: 1389-1391.
- Krishna, S., Maduzia, L.L. and Padgett, R.W. 1999. Specificity of TGF β signaling is conferred by distinct type I receptors and their associated SMAD proteins in *Caenorhabditis elegans*. *Development* 126: 251-260.
- Thatcher, J.D., Haun, C. and Okkema, P.G. 1999. The DAF-3 Smad binds DNA and represses gene expression in the *Caenorhabditis elegans* pharynx. *Development* 126: 97-107.
- Rouault, J.P., Kuwabara, P.E., Sinilnikova, O.M., Duret, L., Thierry-Mieg, D. and Billaud, M. 1999. Regulation of dauer larva development in *Caenorhabditis elegans* by DAF-18, a homologue of the tumour suppressor PTEN. *Curr. Biol.* 9: 329-332.

SOURCE

DAF-3 (cC-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of DAF-3 of *Caenorhabditis elegans* origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9290 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

DAF-3 (cC-19) is recommended for detection of DAF-3 of *Caenorhabditis elegans* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.