HOP-1 (cC-16): sc-9349



The Power to Question

BACKGROUND

Human presenilin proteins have been implicated in the development of Alzheimer's disease. Several presenilin homologs have been identified in *C. elegans*, including SEL-12, HOP-1 and SPE-4. Reducing or eliminating SEL-12 activity causes an egg-laying defective (Egl) phenotype in *C. elegans*. HOP-1 activity can rescue the egg-laying defect of the SEL-12 mutant. SEL-12 and HOP-1 are therefore thought to act as redundant promoters of Notch-pathway signaling by facilitating the activity of LIN-12 and GLP-1. SPE-4 is an integral membrane protein localized within specific organelles responsible for spermatogenesis. Mutation of SPE-4 prevents cytokinesis during meiosis, preventing spermatogenesis.

REFERENCES

- L'Hernault, S.W. and Arduengo, P.M. 1992. Mutation of a putative sperm membrane protein in *Caenorhabditis elegans* prevents sperm differentiation but not its associated meiotic divisions. J. Cell Biol. 119: 55-68.
- Levy-Lahad, E., Wijsman, E.M., Nemens, E., Anderson, L., Goddard, K.A., Weber, J.L., Bird, T.D. and Schellenberg, G.D. 1995. A familial Alzheimer's disease locus on chromosome 1. Science 269: 970-973.
- Sherrington, R., Rogaev, E.I., Liang, Y., Rogaeva, E.A., Levesque, G., Ikeda, M., Chi, H., Lin, C., Li, G., Holman, K., et al. 1995. Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. Nature 375: 754-760.
- 4. Rogaev, E.I., Sherrington, R., Rogaeva, E.A., Levesque, G., Ikeda, M., Liang, Y., Chi, H., Lin, C., Holman, K., Tsuda, T., et al. 1995. Familial Alzheimer's disease in kindreds with missense mutations in a gene on chromosome 1 related to the Alzheimer's disease type 3 gene. Nature 376: 775-778.
- Levitan, D. and Greenwald, I. 1995. Facilitation of LIN-12-mediated signalling by SEL-12, a *Caenorhabditis elegans* S182 Alzheimer's disease gene. Nature 377: 351-354.
- Li, X. and Greenwald, I. 1997. HOP-1, a Caenorhabditis elegans presenilin, appears to be functionally redundant with SEL-12 presenilin and to facilitate LIN-12 and GLP-1 signaling. Proc. Natl. Acad. Sci. USA 94: 12204-12209.

SOURCE

HOP-1 (cC-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of HOP-1 of *Caenorhabditis elegans* origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9349 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

HOP-1 (cC-16) is recommended for detection of HOP-1 of *Caenorhabditis elegans* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

 Rai, R., Zheng, H., He, H., Luo, Y., Multani, A., Carpenter, P.B. and Chang, S. 2010. The function of classical and alternative non-homologous endjoining pathways in the fusion of dysfunctional telomeres. EMBO J. 29: 2598-2610.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com