

# WASP (N-20): sc-9353

## BACKGROUND

The Wiskott-Aldrich syndrome (WAS) is a disorder that results from a monogenic defect that has been mapped to the short arm of the X chromosome at Xp11.23. WAS is characterized by thrombocytopenia, eczema, defects in cell-mediated and humoral immunity and a propensity for lymphoproliferative disease. The gene that is mutated in the syndrome encodes a proline-rich protein of unknown function designated WAS protein (WASP). A clue to WASP function came from the observation that T cells from affected males had an irregular cellular morphology and a disarrayed cytoskeleton suggesting the involvement of WASP in cytoskeletal organization. Close examination of the WASP sequence revealed a putative Cdc42/Rac interacting domain, homologous with those found in PAK65 and ACK. Subsequent investigation has shown WASP to be a true downstream effector of Cdc42.

## CHROMOSOMAL LOCATION

Genetic locus: WAS (human) mapping to Xp11.23; Was (mouse) mapping to X A1.1.

## SOURCE

WASP (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Wiskott-Aldrich syndrome protein (WASP) of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9353 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

WASP (N-20) is recommended for detection of WASP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

WASP (N-20) is also recommended for detection of WASP in additional species, including canine.

Suitable for use as control antibody for WASP siRNA (h): sc-29525, WASP siRNA (m): sc-36830, WASP shRNA Plasmid (h): sc-29525-SH, WASP shRNA Plasmid (m): sc-36830-SH, WASP shRNA (h) Lentiviral Particles: sc-29525-V and WASP shRNA (m) Lentiviral Particles: sc-36830-V.


Molecular Weight of WASP: 66 kDa.

Positive Controls: MOLT-4 cell lysate: sc-2233, BJAB whole cell lysate: sc-2207 or Ramos cell lysate: sc-2216.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



WASP (N-20): sc-9353. Western blot analysis of WASP expression in BJAB (A) and MOLT-4 (B) tissue extracts.

## SELECT PRODUCT CITATIONS

- Lei, Q.Y., Zhang, H., Zhao, B., Zha, Z.Y., Bai, F., Pei, X.H., Zhao, S., Xiong, Y. and Guan, K.L. 2008. TAZ promotes cell proliferation and epithelial-mesenchymal transition and is inhibited by the hippo pathway. *Mol. Cell Biol.* 28: 2426-2436.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **WASP (B-9): sc-13139** or **WASP (F-8): sc-365859**, our highly recommended monoclonal alternatives to WASP (N-20). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **WASP (B-9): sc-13139**.