

CRSP150 (M-19): sc-9419

BACKGROUND

In mammalian cells, transcription is regulated in part by high molecular weight coactivating complexes that mediate signals between transcriptional activators and RNA polymerase. These complexes include CRSP (for cofactor required for Sp1 activation), which is required, in conjunction with TAFII, for transcriptional activation by Sp1. CRSP is ubiquitously expressed in various tissues and functions as a multimeric complex that consists of nine distinct subunits. Several members of the CRSP family share sequence similarity with multiple components of the yeast transcriptional mediator proteins, including CRSP150, which is related to yeast Rgr1, and CRSP70, which is similar to the elongation factor TFIIS. CRSP77 and CRSP150 are also related to proteins within the putative murine mediator complex, while CRSP130 and CRSP34 are largely unrelated to either murine or yeast proteins. CRSP subunits also associate with larger multimeric co-activator complexes, including ARC/DRI, which binds directly to SREBP and nuclear hormone receptors to facilitate transcription, and with NAT, a polymerase II-interacting complex that represses activated transcription.

CHROMOSOMAL LOCATION

Genetic locus: MED14 (human) mapping to Xp11.4.

SOURCE

CRSP150 (M-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of CRSP150 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-9419 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-9419 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CRSP150 (M-19) is recommended for detection of CRSP150 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CRSP150 (M-19) is also recommended for detection of CRSP150 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for CRSP150 siRNA (h): sc-38579, CRSP150 shRNA Plasmid (h): sc-38579-SH and CRSP150 shRNA (h) Lentiviral Particles: sc-38579-V.


CRSP150 (M-19) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Positive Controls: DU 145 nuclear extract: sc-24960, SW480 nuclear extract: sc-2155 or CCRF-CEM nuclear extract: sc-2146.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CRSP150 (M-19): sc-9419. Western blot analysis of CRSP150 expression in A549 (A), DU 145 (B), SW480 (C) and CCRF-CEM (D) nuclear extracts and human heart tissue extract (E).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **CRSP150 (2214C1a): sc-81236**, our highly recommended monoclonal alternative to CRSP150 (M-19).