

Srb7 (V-19): sc-9439

BACKGROUND

In mammalian cells, transcription is regulated in part by high molecular weight coactivating complexes that mediate signals between transcriptional activators and RNA polymerase. These complexes include the SMCC (SRB and MED protein cofactor complex), which consists of various subunits that share homology with several components of the yeast transcriptional mediator complexes, and including the human proteins Srb7, Med6 (also designated DRIP33) and Med7 (also designated DRIP34). SMCC associates with the RNAPII (RNA polymerase II) holoenzyme through Srb7 and, in turn, enhances gene-specific activation or repression induced by DNA-binding transcription factors. Med6 and Med7, as well as other components of SMCC, associate with coactivator proteins from the TRAP (thyroid hormone receptor-activating protein) complex and DRIP (for vitamin D receptor interacting protein) complex to facilitate steroid receptor dependent transcriptional activation. Additionally, SMCC associates with PC4 (positive cofactor 4) to repress basal transcription independent of RNAPII activity.

REFERENCES

1. Malik, S., et al. 1998. A dynamic model for PC4 coactivator function in RNA polymerase II transcription. *Proc. Natl. Acad. Sci. USA* 95: 2192-2197.
2. Jiang, Y.W., et al. 1998. Mammalian mediator of transcriptional regulation and its possible role as an end-point of signal transduction pathways. *Proc. Natl. Acad. Sci. USA* 95: 8538-8543.
3. Gu, W., et al. 1999. A novel human SRB/MED-containing cofactor complex, SMCC, involved in transcription regulation. *Mol. Cell* 3: 97-108.
4. Xiao, H., et al. 1999. The human homologue of *Drosophila* TRF-proximal protein is associated with an RNA polymerase II-SRB complex. *J. Biol. Chem.* 274: 3937-3940.
5. Ito, M., et al. 1999. Identity between TRAP and SMCC complexes indicates novel pathways for the function of nuclear receptors and diverse mammalian activators. *Mol. Cell* 3: 361-370.
6. Rachez, C., et al. 1999. Ligand-dependent transcription activation by nuclear receptors requires the DRIP complex. *Nature* 398: 824-828.

CHROMOSOMAL LOCATION

Genetic locus: SURB7 (human) mapping to 12p11.23.

SOURCE

Srb7 (V-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Srb7 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9439 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-9439 X, 200 µg/0.1 ml.

APPLICATIONS

Srb7 (V-19) is recommended for detection of Srb7 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Srb7 (V-19) is also recommended for detection of Srb7 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Srb7 siRNA (h): sc-38585, Srb7 shRNA Plasmid (h): sc-38585-SH and Srb7 shRNA (h) Lentiviral Particles: sc-38585-V.

Srb7 (V-19) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of Srb7: 16 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.