

PA26-T3 (L-18): sc-9508

BACKGROUND

Cell cycle progression is subject to arrest at G₁ and G₂ checkpoints in response to DNA damage, presumably to allow time for DNA repair prior to entry into S and M phase, respectively. The p53 tumor suppressor is required for one such G₁ checkpoint and functions to upregulate expression of GADD 45 and the mitotic inhibitory protein p21. GADD 45 stimulates DNA excision repair *in vitro* and inhibits entry of cells into S phase, and it apparently acts in concert with GADD 153 in inducing growth arrest. A related DNA-damage inducible gene, GADD 34 synergizes with GADD 45 or GADD 153 in suppressing cell growth. PEG-3 (progression elevated gene-3) shares significant homology with GADD 34 and is inducible by DNA damage. An additional GADD related gene, PA26, is a possible target of p53. Three isoforms of PA26 have been identified as PA26-T1, PA26-T2 and PA26-T3.

REFERENCES

1. Sherr, C.J. 1994. G₁ phase progression: cycling on cue. *Cell* 79: 551-555.
2. Hunter, T. et al. 1994. Cyclins and cancer II: cyclin D and CDK inhibitors come of age. *Cell* 79: 573-582.

CHROMOSOMAL LOCATION

Genetic locus: SESN1 (human) mapping to 6q21; Sesn1 (mouse) mapping to 10 B2.

SOURCE

PA26-T3 (L-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PA26-T3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9508 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PA26-T3 (L-18) is recommended for detection of all PA26-T isoforms of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PA26-T3 (L-18) is also recommended for detection of all PA26-T isoforms in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for PA26 siRNA (h): sc-37420, PA26 siRNA (m): sc-37421, PA26 shRNA Plasmid (h): sc-37420-SH, PA26 shRNA Plasmid (m): sc-37421-SH, PA26 shRNA (h) Lentiviral Particles: sc-37420-V and PA26 shRNA (m) Lentiviral Particles: sc-37421-V.


Molecular Weight of PA26-T3: 57 kDa.

Positive Controls: PA26 (m): 293T Lysate: sc-122341.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



PA26-T3 (L-18): sc-9508. Western blot analysis of PA26 expression in non-transfected: sc-117752 (A) and mouse PA26 transfected: sc-122341 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **PA26 (C-10): sc-376170**, our highly recommended monoclonal alternative to PA26-T3 (L-18).