

cyclin I (D-19): sc-9534



The Power to Question

BACKGROUND

Cyclins are the regulatory subunits of Cdc2 p34 and related cyclin-dependent kinases (Cdks) which play critical roles in the control of cell cycle progression. The catalytic subunit for cyclin A and B is Cdc2 p34 kinase. The Cdc2-cyclin B complex controls the G₂ to M transition whereas Cdc2-cyclin A regulates S phase progression. Cyclin D1 accumulates during G₁ and associates with Cdk2, Cdk4 and Cdk5. Cyclin E and Cdk2 interact during the G₁ to S transition. Cyclin G contains a typical N terminal cyclin box and a carboxy terminal domain sequence homologous to the tyrosine phosphorylation site of the epidermal growth factor receptor. Cyclin G₂ shares 53% amino acid sequence identity with cyclin G₁. Cyclin I shares highest sequence similarity to cyclins G and E and is most highly expressed in skeletal muscle, heart and brain.

REFERENCES

1. Pines, J., et al. 1990. Human cyclin A is adenovirus E1A-associated protein p60 and behaves differently from cyclin B. *Nature* 346: 760-763.
2. Fang, F., et al. 1991. Evidence that the G₁-S and G₂-M transitions are controlled by different cdc2 proteins in higher eukaryotes. *Cell* 66: 731-742.
3. Koff, A., et al. 1991. Human cyclin E, a new cyclin that interacts with two members of the CDC2 gene family. *Cell* 66: 1217-1228.
4. Girard, F., et al. 1991. Cyclin A is required for the onset of DNA replication in mammalian fibroblasts. *Cell* 67: 1169-1179.
5. Xiong, Y., et al. 1992. D type cyclins associate with multiple protein kinases and the DNA replication and repair factor PCNA. *Cell* 71: 505-514.
6. Tamura, K., et al. 1993. Cyclin G: a new mammalian cyclin with homology to fission yeast Cig1. *Oncogene* 8: 2113-2118.
7. Nakamura, T., et al. 1995. Cyclin I: a new mcyclin encoded by a gene isolated from human brain. *Exp. Cell Res.* 221: 534-542.
8. Horne, M.C., et al. 1996. Cyclin G₁ and cyclin G₂ comprise a new family of cyclins with contrasting tissue-specific and cell cycle-regulated expressions. *J. Biol. Chem.* 271: 6050-6061.

CHROMOSOMAL LOCATION

Genetic locus: CCNI (human) mapping to 4q21.1; Ccni (mouse) mapping to 5 E2.

SOURCE

cyclin I (D-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of cyclin I of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9534 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

cyclin I (D-19) is recommended for detection of cyclin I of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

cyclin I (D-19) is also recommended for detection of cyclin I in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for cyclin I siRNA (h): sc-35141, cyclin I siRNA (m): sc-35142, cyclin I shRNA Plasmid (h): sc-35141-SH, cyclin I shRNA Plasmid (m): sc-35142-SH, cyclin I shRNA (h) Lentiviral Particles: sc-35141-V and cyclin I shRNA (m) Lentiviral Particles: sc-35142-V.

Molecular Weight of cyclin I: 47 kDa.

Positive Controls: Rat skeletal muscle extract or mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.