

# caspase-14 p10 (C-20): sc-9644

## BACKGROUND

A unique family of cysteine proteases has been described that differs in sequence, structure and substrate specificity from any previously described protease family. This family, termed Ced-3/caspase-1, is composed of caspase-1, caspase-2, caspase-3, caspase-4, caspase-6 and caspase-7 (also designated Mch3, ICE-LAP3 or CMH-1), caspase-9, caspase-10 and caspase-14. Ced-3/caspase-1 family members function as key components of the apoptotic machinery and act to destroy specific target proteins which are critical to cellular longevity. caspase-3, caspase-7 and caspase-9, but not caspase-1, have been shown to cleave the nuclear protein PARP into an apoptotic fragment. caspase-14, also designated MICE (for mini-ICE), is highly expressed in embryonic tissues but appears to be absent from adult tissues. Procaspase-14 can be processed *in vitro* by caspase-8 and caspase-10 but not by other caspases.

## REFERENCES

1. Duan, H., et al. 1996. ICE-LAP3, a novel mammalian homologue of the *Caenorhabditis elegans* cell death protein Ced-3 is activated during FAS- and tumor necrosis factor-induced apoptosis. J. Biol. Chem. 271: 1621-1625.
2. Duan, H., et al. 1996. ICE-LAP6, a novel member of the ICE/Ced-3 gene family, is activated by the cytotoxic T cell protease granzyme B. J. Biol. Chem. 271: 16720-16724.
3. Fernandes-Alnemri, T.F., et al. 1996. *In vitro* activation of CPP32 and Mch3 by Mch4, a novel human apoptotic cysteine protease containing two FADD-like domains. Proc. Natl. Acad. Sci. USA 93: 7464-7469.
4. Casciola-Rosen, L., et al. 1996. Apopain/CPP32 cleaves proteins that are essential for cellular repair: a fundamental principle of apoptotic death. J. Exp. Med. 183: 1957-1964.

## CHROMOSOMAL LOCATION

Genetic locus: CASP14 (human) mapping to 19p13.12.

## SOURCE

caspase-14 p10 (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of caspase-14 p10 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9644 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

caspase-14 p10 (C-20) is recommended for detection of the p10 subunit and precursor of caspase-14 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


caspase-14 p10 (C-20) is also recommended for detection of the p10 subunit and precursor of caspase-14 in additional species, including equine, canine, bovine and porcine.

Molecular Weight of caspase-14 p10: 30/18/11 kDa.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



caspase-14 p10 (C-20): sc-9644. Western blot analysis of human recombinant procaspase-14.



caspase-14 p10 (C-20): sc-9644. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skin tissue showing cytoplasmic staining of epidermal cells, keratinocytes, Langerhans cells and melanocytes.

## SELECT PRODUCT CITATIONS

1. Hibino, T., et al. 2010. Purification and characterization of active caspase-14 from human epidermis and development of the cleavage site-directed antibody. J. Cell. Biochem. 109: 487-497.

**MONOS**  
Satisfaction  
Guaranteed

Try **caspase-14 (D-10): sc-48336** or **caspase-14 (C-12): sc-48395**, our highly recommended monoclonal alternatives to caspase-14 p10 (C-20).