

# ZNF845 siRNA (h): sc-97085

## BACKGROUND

ZNF845 (zinc finger protein 845) is a 970 amino acid protein that belongs to the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc-finger protein family. Members of this family have been linked to autosomal recessive congenital microcephaly (Jawad syndrome), human colorectal cancer and myocardial infarction via transcriptional regulation. ZNF845 contains 27 C<sub>2</sub>H<sub>2</sub>-type zinc fingers as well as 1 KRAB domain, and may be involved in transcriptional regulation. ZNF845 localizes to nucleus and is encoded by a gene that maps to chromosome 19q13.42. Chromosome 19 consists of over 63 million bases, houses approximately 1,400 genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin (Ig) superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc receptors (FcRs).

## REFERENCES

1. Ashworth, L.K., Batzer, M.A., Brandriff, B., Branscomb, E., de Jong, P., Garcia, E., Garnes, J.A., Gordon, L.A., Lamerdin, J.E., Lennon, G., Mohrenweiser, H., Olsen, A.S., Slezak, T. and Carrano, A.V. 1995. An integrated metric physical map of human chromosome 19. *Nat. Genet.* 11: 422-427.
2. Poloumienko, A. 2004. Cloning and comparative analysis of the bovine, porcine, and equine sex chromosome genes ZFX and ZFY. *Genome* 47: 74-83.
3. Tian, X., Sun, D., Zhang, Y., Zhao, S., Xiong, H. and Fang, J. 2008. Zinc finger protein 278, a potential oncogene in human colorectal cancer. *Acta Biochim. Biophys. Sin.* 40: 289-296.
4. Hassan, M.J., Chishti, M.S., Jamal, S.M., Tariq, M. and Ahmad, W. 2008. A syndromic form of autosomal recessive congenital microcephaly (Jawad syndrome) maps to chromosome 18p11.22-q11.2. *Hum. Genet.* 123: 77-82.
5. Yamada, Y., Kato, K., Oguri, M., Fujimaki, T., Yokoi, K., Matsuo, H., Watanabe, S., Metoki, N., Yoshida, H., Satoh, K., Ichihara, S., Aoyagi, Y., Yasunaga, A., Park, H., Tanaka, M. and Nozawa, Y. 2008. Genetic risk for myocardial infarction determined by polymorphisms of candidate genes in a Japanese population. *J. Med. Genet.* 45: 216-221.
6. Alfonso, P., Cañamero, M., Fernández-Carbonie, F., Núñez, A. and Casal, J.I. 2008. Proteome analysis of membrane fractions in colorectal carcinomas by using 2D-DIGE saturation labeling. *J. Proteome Res.* 7: 4247-4255.
7. Kumar, V., Hassan, M.I., Kashav, T., Singh, T.P. and Yadav, S. 2008. Heparin-binding proteins of human seminal plasma: purification and characterization. *Mol. Reprod. Dev.* 75: 1767-1774.
8. SWISS-PROT/TrEMBL (Q96IR2). World Wide Web URL: <http://www.uniprot.org/uniprot/Q96IR2>

## CHROMOSOMAL LOCATION

Genetic locus: ZNF845 (human) mapping to 19q13.42.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

ZNF845 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF845 shRNA Plasmid (h): sc-97085-SH and ZNF845 shRNA (h) Lentiviral Particles: sc-97085-V as alternate gene silencing products.

For independent verification of ZNF845 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-97085A, sc-97085B and sc-97085C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ZNF845 siRNA (h) is recommended for the inhibition of ZNF845 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF845 gene expression knockdown using RT-PCR Primer: ZNF845 (h)-PR: sc-97085-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.