

AbdA (d-130): sc-98261

BACKGROUND

Drosophila melanogaster, a proven and effective model for studying developmental and cellular processes common to higher eukaryotes, contains a genome encoding approximately 13,600 genes, which were elucidated from more than 120 megabases of euchromatin. These genes are organized among chromosomes 2, 3, 4, X and Y, with the Y chromosome being predominately heterochromatic. *Drosophila* genes, which are categorized based on the type of protein for which they encode, represent six major classifications, including intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing and chromatin associated) and other functional proteins. Morphogenesis and cell differentiation in *Drosophila* requires accurate control of cell division. The Hox/homeotic genes encode transcription factors that generate segmental diversity during *Drosophila* development. In *Drosophila*, the Ultrabithorax (Ubx) and abdominal A (AbdA, also abd-A) Hox proteins are expressed largely in the abdominal segments, where they suppress thoracic leg development during embryogenesis.

REFERENCES

1. Lehner, C.F. 1991. Pulling the string: cell cycle regulation during *Drosophila* development. *Semin. Cell Biol.* 2: 223-231.
2. Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P., et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185-2195.
3. Mata, J., Curado, S., Ephrussi, A. and Rorth, P. 2000. Tribbles coordinates mitosis and morphogenesis in *Drosophila* by regulating string/CDC25 proteolysis. *Cell* 101: 511-522.
4. Brodu, V., Elstob, P.R. and Gould, A.P. 2002. Abdominal A specifies one cell type in *Drosophila* by regulating one principal target gene. *Development* 129: 2957-2963.
5. Ronshaugen, M., McGinnis, N. and McGinnis, W. 2002. Hox protein mutation and macroevolution of the insect body plan. *Nature* 415: 914-917.

SOURCE

AbdA (d-130) is a rabbit polyclonal antibody raised against amino acids 261-390 mapping within an internal region of AbdA of *Drosophila melanogaster* origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

AbdA (d-130) is recommended for detection of AbdA of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **AbdA (C-11): sc-390990**, our highly recommended monoclonal alternative to AbdA (d-130).