

FMR2 (H-185): sc-98361

BACKGROUND

FMR2 (fragile X mental retardation 2), also known as AF4 family member 2 (AFF2) or protein Ox19, is a 1,311 amino acid nuclear protein belonging to the AF4 family. Expressed in the brain, placenta and lung, FMR2 exists as two isoforms produced by alternative splicing. Defects in the gene that encodes FMR2 have been found to be a cause of FRAXE, an X-linked form of mental retardation. Individuals expressing the FRAXE site also have more than 200 copies of a GCC repeat adjacent to CpG island, compared to 6 to 35 copies of the GCC repeat in a normal individual. It is believed that loss of FMR2 expression causes this GCC expansion of the FRAXE site.

REFERENCES

1. Geçz, J., Bielby, S., Sutherland, G.R. and Mulley, J.C. 1997. Gene structure and subcellular localization of FMR2, a member of a new family of putative transcription activators. *Genomics* 44: 201-213.
2. Geçz, J., Oostra, B.A., Hockey, A., Carbonell, P., Turner, G., Haan, E.A., Sutherland, G.R. and Mulley, J.C. 1997. FMR2 expression in families with FRAXE mental retardation. *Hum. Mol. Genet.* 6: 435-441.
3. Chakrabarti, L., Bristulf, J., Foss, G.S. and Davies, K.E. 1998. Expression of the murine homologue of FMR2 in mouse brain and during development. *Hum. Mol. Genet.* 7: 441-448.
4. Geçz, J. and Mulley, J.C. 1999. Characterisation and expression of a large, 13.7 kb FMR2 isoform. *Eur. J. Hum. Genet.* 7: 157-162.
5. Sharma, D., Gupta, M. and Thelma, B.K. 2001. Expansion mutation frequency and CGG/GCC repeat polymorphism in FMR1 and FMR2 genes in an Indian population. *Genet. Epidemiol.* 20: 129-144.
6. Hillman, M.A. and Geçz, J. 2001. Fragile XE-associated familial mental retardation protein 2 (FMR2) acts as a potent transcription activator. *J. Hum. Genet.* 46: 251-259.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 309548. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Brylawski, B.P., Chastain, P.D., Cohen, S.M., Cordeiro-Stone, M. and Kaufman, D.G. 2007. Mapping of an origin of DNA replication in the promoter of fragile X gene FMR1. *Exp. Mol. Pathol.* 82: 190-196.

CHROMOSOMAL LOCATION

Genetic locus: AFF2 (human) mapping to Xq28; Aff2 (mouse) mapping to X A7.1.

SOURCE

FMR2 (H-185) is a rabbit polyclonal antibody raised against amino acids 81-265 mapping near the N-terminus of FMR2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

FMR2 (H-185) is recommended for detection of FMR2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FMR2 siRNA (h): sc-75046, FMR2 siRNA (m): sc-75047, FMR2 shRNA Plasmid (h): sc-75046-SH, FMR2 shRNA Plasmid (m): sc-75047-SH, FMR2 shRNA (h) Lentiviral Particles: sc-75046-V and FMR2 shRNA (m) Lentiviral Particles: sc-75047-V.


Molecular Weight of FMR2: 140 kDa.

Positive Controls: OVCAR-3 whole cell lysate.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



FMR2 (H-185): sc-98361. Western blot analysis of FMR2 expression in OVCAR-3 whole cell lysate.



FMR2 (H-185): sc-98361. Immunofluorescence staining of formalin-fixed HeLa cells showing nuclear and cytoplasmic localization. Kindly provided by Yang Xiang, Ph.D., Division of Newborn Medicine, Boston Children's Hospital, Cell Biology Department, Harvard Medical School.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.